



# Journal of Pharmaceutical Research and Reviews (JPRR)



## Medicinal plants used to treat the disease of gout –an overview

R.Ramasubramania raja M.Sekar, P.Lathasri,I.Paula Preethi,M.Suma Bhavana,T.Swathi

Department of Pharmacognosy, Narayana Pharmacy College, Nellore, A.P

### ABSTRACT

Gout also called as hyper uricemia. Gout is characterized by sudden, severe attacks of pain, redness and tenderness in joints, often the joint at the base of big toe. Gout is a condition that develops when Uric acid is not properly excreted from the blood through urine .Excess uric acid builds up in the body and when not removed by the kidneys, it deposits in the form of crystals in the joints, primarily in hands and feet and especially in the joint of big toe. The risk of developing gout increases as the Serum uric acid level rises [ $>7$  mg/dl]. In our focus medicinal plants used to treat the gout for the reason in the allopathic system of medicine is not cure the disease of gout, its only controlled. So few medicinal plants we reviewed this paper.

**Keywords:** Gout, uricemia, pain, serum, medicinal plants

### \*Correspondence to Author:

Dr.R.Ramasubramania raja  
Department of Pharmacognosy,  
Narayana Pharmacy College, Nel-  
lore, A.P

### How to cite this article:

R.Ramasubramania raja M. Sekar,  
P.Lathasri,I.Paula Preethi,M.Suma  
Bhavana,T.Swathi Medicinal plants  
used to treat the disease of gout  
–an overview. Journal of Pharma-  
ceutical Research and Reviews,  
2018; 2:6.

**eSciencePublisher**

eSciPub LLC, Houston, TX USA.

Website: <http://escipub.com/>

## Introduction

Gout disease, first identified by Egyptians in 2640 BC, PODAGRA [Acute gout occurring in the first Metatarsophalangeal joint] was later recognized by Hippocrates in 15<sup>th</sup> century BC who referred to it as the Un walkable disease throughout history, Gout has been associated with Rich foods and excessive alcohol consumption. There is evidence that Colchicine, an alkaloid derived from the Autumn crocus [*Colchicum autumnale*] was used as powerful purgative in ancient Greece, its first use as a selective and specific treatment for Gout. Uricosuric agents were first used at the end of 19 centuries. In the modern era Non steroidal anti-inflammatory drugs are usually the drugs of choice for treating acute gout. Xanthine oxidase inhibitors, which are effective in reducing plasma and urinary urate levels.

## Definition:

Gout also called as hyper uricemia. Gout is characterized by sudden, severe attacks of pain, redness and tenderness in joints, often the joint at the base of big toe. Gout a complex form of Arthritis Gout can affect any one, Men are more likely to be affected than Women. become increasingly susceptible to get gout after "Menopause.



**Fig: showing the inflammation in joints due to uric acid accumulation**

An attack of gout can occur suddenly, often waking you up in the middle of night with the sensation that big toe is on fire. The affected joint is hot, swollen and so tender that even the weight of the sheet on it may be intolerable.

Gout is a condition that develops when Uric acid is not properly excreted from the blood

through urine. Excess uric acid builds up in the body and when not removed by the kidneys, it deposits in the form of crystals in the joints, primarily in hands and feet and especially in the joint of big toe. Gout a very painful form of Arthritis, seen mostly in men over 50 years of age. Now-a-days it is seen in the wider range of populations, as well as in women and children and is thought to be the result of eating habits where people consume excess of fats which contribute heavily to the formation of purine break down and uric acid. Uric acid is deposited in the form of "Monosodium urate crystals" in the various tissues. [For eg...Joints, Connective tissue, Kidney]

**PRIMARY HYPERURICEMIA:** Is called when Uric acid crystals saturation arises without coexisting diseases or drugs that alter uric acid production and secretion. In **SECONDARY HYPER URICEMIA** there is an excessive uric acid production or diminished renal clearance that occurs as a result of disease drug dietary or toxin.

## SIGNS & SYMPTOMS:

The signs and symptoms of gout almost always occur suddenly often at night and without warning. They include: Intense joint pain, Lingering discomfort, [after most severe pain subsides, some joint discomfort may last from few days to few weeks]. Inflammation and redness limited range of motion. Acute and chronic inflammation associated with changes in Articular and periarticular structures. The risk of developing gout increases as the Serum uric acid level rises [ $>7$  mg/dl]

## FACTORS THAT INCREASING THE URIC ACID IN THE BODY INCLUDE:

**DIET:** Eating a diet that's high in meat and seafood and high in beverages sweetened with fruit sugar [fructose] promotes higher uric acid levels, which increases risk of Gout. Alcohol consumption, especially of beer, also increases the risk of Gout.

**OBESITY:** If overweight, body produces more Uric acid and kidneys have more difficulty in eliminating uric acid which greatly increases the risk of Gout.

**MEDICAL CONDITIONS:**

Certain diseases and conditions also develop Gout. These conditions include: untreated high blood pressure and chronic conditions like Diabetes, Metabolic syndrome, Heart and Kidney diseases. Hematological causes of hyperuricemia and gout associated with increased nucleic acid turnover and break down to uric acid. Lymphoproliferative disorders, myeloproliferative disorders, certain hemolytic anemias and hemoglobinopathies. In chronic renal failure condition reduced clearance of uric acid can lead to Hyperuricemia.

**CERTAIN MEDICATIONS:**


The use of THIAZIDE DIURETICS [commonly used to treat hypertension] and low dose of ASPIRIN also can increase uric acid levels. CYTOTOXIC DRUGS increase uric acid concentrations by enhancing nucleic acid turnover and excretion. ETHAMBUTANOL and NICOTINIC ACID increase the uric acid concentrations by competing with urate for tubular secretion sites, thereby decreasing uric acid excretion. CYCLOSPORINE decreases renal urate clearance as Pyrazinamide and Levodopa. Miscellaneous disorders like Keto acidosis, Psoriasis, Chronic lead poisoning are examples of conditions that may cause Hyperuricemia.

**FAMILY HISTORY OF GOUT:** If other members of family have had Gout, then the persons of that family are more likely to develop the gout disease.

**RECENT SURGERY OR TRAUMA:**


Experiencing recent surgery or trauma has been associated with increased risk of developing Gout. **AGE & SEX:** Gout occurs more often in men, primarily because, women tend to have lower uric acid levels. After “menopause”, women’s uric acid levels approach those of men. Men are more likely to develop gout earlier usually between 30 and 50 years of age, whereas Women generally develop signs and symptoms after menopause.

**MECHANISM OF GOUT DISEASE**



**Fig: formation of inflammation due to accumulation of uric acid**

## ALLOPATHIC MEDICINE TO TREAT GOUT:



**Anti inflammatory** drug therapy should begin immediately for maximal therapeutic effectiveness. **Urate lowering drugs** should not be given until the acute attack is controlled. As these drugs may prolong the attack by causing a change in Uric acid equilibrium. Urate lowering agents can begin within 1 to 2 weeks after resolution of the attack. Specific drugs given in Gout are: **NSAIDS [Non steroidal anti inflammatory drugs ]**: These drugs preferred when treatment is delayed significantly after symptoms onset or when the patient cannot tolerate the adverse GI effects of Colchicine.


**Indomethacin [Indocin]** : Usually given in dose of 50 mg three times daily initially then the dose is tapered. **NAPROXEN [Naprosyn ] : 750 mg** followed by 250 mg every 8 hours until the attack subsides. **SULINDAC [Clinoril]**: 200mg twice day to short reduce the dose with satisfactory response. Some times Cyclo oxygenase 2 [COX-2] inhibitors also may be used for these attacks.

## CORTICOSTEROIDS:

**Corticosteroid Intra articular injections** are effective in patients with acute single joint gout. Specific doses of corticosteroids are related to joint size. An intra articular dose of **Triamcinolone** -8mg in smaller joints [10 mg in the knee], **Methylprednisolone acetate** -5 to 25 mg per joints, **Betamethasone** – 3 to 6 mg can be used.

**Systemic Corticosteroid Therapy:** It is especially used when either NSAIDS or Colchicine cannot be given or have not been effective. Oral Prednisone – 20 to 60 mg /day initially with the dose tapered during a period of 5 to 7 days. It can be useful for gout patients without producing the rebound effect. **IM Betamethasone** -7mg or **IV Methyl prednisolone** – 125 mg are examples of other potential agents. **IM ACTH [Adrenocorticotropin hormone ]** – 40 to 80 units helps in reducing inflammation by stimulating corticosteroid production by the patients Adrenal gland plus it activates Melanocortin type 3 receptors. These agents cannot be preferred because of short duration of action.

## COLCHICINE:



**Fig : Showing the effects of Colchicine in various body parts**

It is an Antimitotic drug, highly effective at relieving acute attacks of Gout but has the lowest benefit or toxicity. A typical oral dose is 0.5 to 0.6 mg every 1 to 2 hours until the pain is abated. Maximum dose – 6 to 8 mg has been administered.

**ALLOPURINOL:** Dose – 100 to 300 mg per day orally. Currently it is most commonly used agents for long term control of Chronic Gout. It is the drug of **Xanthine oxidase** enzyme that catalyses the conversion of Xanthine to uric acid and Hypoxanthine to Xanthine. Major active metabolite in it is Oxypurinol also a Xanthine oxidase inhibitor.

**URICOSURIC AGENTS:** It is generally recommended for younger patients with normal



renal function who are under excretors of uric acid . Both Probenecid and Sulfinpyrazone are competitively inhibit the active reabsorption of uric acid at PCT . It helps in decrease the serum urate concentrations .In **presence of renal dysfunction [creatinine clearance <50 ml/min ] uricosurics are not effective and are contraindicated .**

## HERBAL MEDICINE FOR TREATMENT OF GOUT:

Healthy food choices that can help in regulating uric acid in the body.

### GREEN TEA:



**Fig: Green tea**

Scientific name: *Camellia sinensis*

Biological source : *Camellia sinensis* is a species evergreen shrub or small tree whose leaves and leaf buds are used to produce Tea

Family: Theaceae

Active constituents : Catechins, Caffeine, and Theanine (amino acid) Uses : Antioxidant, Antibacterial Green tea is rich with high Catechin content which is a potent antioxidant. Catechin helps in slow down the production of particular type of enzymes in the body that is related to uric acid formation. Thus it extremely helps in reducing the uric acid levels.

**FIBRE:** Fibre rich diet can be helpful in reduces the uric acid level in the body.

Examples..... Oats, whole grains

### OATS



**Fig: Oats**

Scientific name : *Avena sativa*

Family: Poaceae

Active constituents: Avenanthramides, Flavavanoligans , Beta glucan ,Terpenoid  
Uses:Reduce the uric acid level in the body

Vegetables like Broccoli, Pumpkin, Celery also helps in reducing the uric acid level, these type of foods filled with dietary fibres that are extremely beneficial in absorption of uric acid and eliminating it from the body.

### BROCCOLI



**Fig: Broccoli**

Scientific name : *Brassica oleracea*

Biological source: species of *Brassica oleracea*

Family: cruciferae

Active constituents: Provitamin A, Vitamin C ,Vitamin K, Isothiocyanates

Uses : Reduces the uric acid levels in the body

### PUMPKIN



**Fig: cucurbits**

Scientific name:*Cucurbita maxima*

Family : Cucurbitaceae

Biological source: It is cultivated in the form of *Cucurbita maxima* , *C. moschata*, *C.argyrosperma*

Active constituents:Phytosterols , (eg... Betasitosterol)

### CELERY

Scientific name: *Apium graveolens*

Family : Apiaceae /Umbelliferae

Biological source : Long fibrous stalk tapering into leaves of Celery Active constituents :

Hydroxyphenyl ethanol, Dimethoxy psoralen  
, Isofraxidin ,Eugenic acid  
Uses: Spice ,and its extracts have been used  
as Herbal medicine

**VITAMIN C ENRICHED FRUITS:**

These reduce the uric acid level in short period  
.The required amount of vitamin c every day is  
500 mg

Examples: Orange, Sweet lemon

**ORANGE:**



**Fig: Orange**

Scientific name: Citrus sinensis

Family: Rutaceae

Active constituents: Hesperidin, Limonene,  
Naringin, Tangaretin, Synephrine alkaloids

**SWEET LEMON:**



**Fig: Sweet lemon**

Scientific name: Citrus limetta

Family: Rutaceae

Biological source: The fruits of Citrus limonum  
species

Active constituents: Vitamin C, Dietary fibre,  
Carbohydrates

**CHERRIES**



**Fig: Cherries**

Scientific name: Prunus avium (sweet cherry)  
,Prunus cerasus (Sourcherry)

Family: Rosaceae

Active constituents: Anthocyanins,  
Hydroxycinnamic acids, Hydroxy benzoic acid  
Uses:It helps in people who suffering with high  
levels of uric acid in their body

**BERRIES:** All types of berries like Straw berry,  
Blue berry, Rasp berry has anti inflammatory  
properties and also contain high in fibre content  
help in promoting reduction of Uric acid in their  
blood .

**BLUE BERRY:**



**Fig: Blue berry**

Scientific name :Cyanococcus

Family :Ericaceae

Active constituents :Pterostilbene, Natural  
antioxidants

Uses:Antioxidants protect us from disease and  
age related health risks .

**RASP BERRY:**



**Fig: Rasp berry**

Scientific name :Rubusidaeus

Family : Rosaceae

Active constituents : Anticholine esterase

Use:Act as smooth muscle stimulant

**STRAW BERRY**



**Fig : Strawberry**

Scientific name: Fragaria ,

Family:Rosaceae

Active constituents : Butanoic acid , Methyl  
ester , Hexanoic acid ethyl ester

Berries have anti inflammatory properties made  
out of substance called Anthocyanins .This  
particular substance helpful in reducing high  
uric acid levels .It also prevents the uric acid  
from crystallizing and getting deposited in the  
joints that may later lead to joint pains.

## FRUITS & TOMATOES



**Fig: Solanum lycopersicum**

### (Tomato)

Fruits, just like vegetables extremely helpful in providing relief from increasing levels of Uric acid. Tomatoes which are also counted as a fruit rather than vegetables and are good for body and their high vitamin -c content can help in reducing the uric acid levels.

Scientific name: Solanum lycopersicum

Family: Solanaceae

Active constituents: Steroids, Proteins, amino acids, Minerals, lipids

### WATER (OR) FLUIDS

Water is one natural cleanser and fluid that flushes out toxins from the body. So definitely drink at least 10 to 12 glasses of water daily .Fluids are beneficial in helping removal of Uric acid from the body through excretion . Drinking more fluids and water help in reducing the uric acid levels.**But remember to choose the right type of juices and fluids.**

### Conclusion:

Gout a very painful form of Arthritis, seen mostly in men over 50 years of age. Now-a-days it is seen in the wider range of populations, as well as in women and children and is thought to be the result of eating habits where people consume excess of fats which contribute heavily to the formation of purine break down and uric acid. India is a developing country, here poor people is more affected this type of the diseases. Without side effect the disease of gout cure for medicinal herbs. In the above mentioning the medicinal herbs once total work of research over it will be help for our society.

### References:

- 1) Comprehensive Pharmacy Review by Leon shargel, Alan H Mutnick
- 2) Woolliscroft JO, Colfer H, Fox IH. Hyperuricemia in acute illness: a poor prognostic sign. Am J Med. 1982;72:58-62. [PMID: 7058824]
- 3) Mineo I, Kono N, Hara N, Shimizu T, Yamada Y, Kawachi M, et al.
- 4) Lieber CS, Jones DP, Losowsky MS, Davidson CS. I
- 5) Gibson T, Rodgers AV, Simmonds HA, Toseland P. 1984;23:203-9.[PMID 6743968]
- 6) Glynn RJ, Campion EW, Silbert JE. 1983;26:87-93. [PMID: 6824508]
- 7) Freedman DS, Khan LK, Serdula MK, Galuska DA, Dietz WH. JAMA. 2002;288:1758-61. [PMID: 12365960]
- 8) Rheum Dis. 2008 Jul;67(7):960–6. [PMC free article] [PubMed] Zhu Y, Pandya BJ, Choi HK.
- 9) Thompson GR, Duff IF, Robinson WD, et al. Long term uricosuric therapy in gout. Arthritis Rheum. 1962 Aug;5:384–96. [PubMed].
- 10) NHANES 2007-2008. Am J Med. 2012 Jul;125(7):679–87 e1. [PubMed].
- 11) Drug interactions by CIMS India.
- 12) Disease out line by disease dictionary.

