



American Journal of Biotechnology and Bioscience (AJBB)



Different statistical methods to collect information about the adverse effects of climatic factors on cotton production

Zakaria M. Sawan

Cotton Research Institute, Agricultural Research Center, Ministry of Agriculture & Land Reclamation, 9 Gamaa Street, 12619, Giza, Egypt

ABSTRACT

This study investigates the statistical relationship between various climatic factors and overall flower and boll production. Also, predicting effects of climatic factors during different convenient intervals (in days) on cotton flower and boll production compared with daily observations. Further, collects information about the nature of the relationship between various climatic factors and cotton boll development and the 15-day period both prior to and after initiation of individual bolls. And, provide information on the effect of various climatic factors and soil moisture status during the development stage on flower and boll production in cotton. Evaporation, sunshine duration, relative humidity, surface soil temperature at 1800 h, and maximum air temperature, are the important climatic factors that significantly affect flower and boll production. The five-day interval was found to be more adequately and sensibly related to yield parameters. Evaporation; minimum humidity and sunshine duration were the most effective climatic factors during preceding and succeeding periods on boll production and retention. There was a negative correlation between flower and boll production and either evaporation or sunshine duration, while that correlation with minimum relative humidity was positive.

Keywords: cotton flower and boll production, evaporation, relative humidity, soil moisture status, sunshine duration, temperature

*Correspondence to Author:

Zakaria M. Sawan
Cotton Research Institute, Agricultural Research Center, Ministry of Agriculture & Land Reclamation, 9 Gamaa Street, 12619, Giza, Egypt (zmsawan@hotmail.com)

How to cite this article:

Zakaria M. Sawan. Different statistical methods to collect information about the adverse effects of climatic factors on cotton production. American Journal of Biotechnology and Bioscience, 2017; 1:1.

eSciencePublisher®

eSciPub LLC, Houston, TX USA.
Website: <http://escipub.com/>

1. Introduction

Climate affects crop growth interactively, sometimes resulting in unexpected responses to prevailing conditions. Many factors, such as length of the growing season, climate (including solar radiation, temperature, light, wind, rainfall, and dew), cultivar, availability of nutrients and soil moisture, pests and cultural practices affect cotton growth (El-Zik 1980). The balance between vegetative and reproductive development can be influenced by soil fertility, soil moisture, cloudy weather, spacing and perhaps other factors such as temperature and relative humidity (Guinn 1982). Weather, soil, cultivars, and cultural practices affect crop growth interactively, sometimes resulting in plants responding in unexpected ways to their conditions (Hodges et al. 1993).

Water is a primary factor controlling plant growth. Xiao et al. (2000) stated that, when water was applied at 0.85, 0.70, 0.55 or 0.40 ET (evapotranspiration) to cotton plants grown in pots, there was a close relationship between plant development and water supply. The fruit-bearing branches, square and boll numbers and boll size were increased with increased water supply. Barbour and Farquhar (2000) reported on greenhouse pot trials where cotton cv. CS50 plants were grown at 43 or 76% relative humidity (RH) and sprayed daily with abscisic acid (ABA) or distilled water. Plants grown at lower RH had higher transpiration rates, lower leaf temperatures and lower stomatal conductance. Plant biomass was also reduced at the lower RH. Within each RH environment, increasing ABA concentration generally reduced stomatal conductance, evaporation rates, superficial leaf density and plant biomass, and increased leaf temperature and specific leaf area.

Temperature is also a primary factor controlling rates of plant growth and development. Burke et al. (1988) has defined the optimum temperature range for biochemical and metabolic activities of plants as the thermal kinetic window (TKW). Plant temperatures above or below the TKW result in stress that limits growth and yield. The TKW for cotton growth is 23.5 to 32°C, with an optimum temperature of 28°C. Biomass production is directly related to the amount of time that foliage temperature is within the TKW. Hodges et al.

(1993) found that the optimum temperature for cotton stem and leaf growth, seedling development, and fruiting was almost 30°C, with fruit retention decreasing rapidly as the time of exposure to 40°C increased. Reddy et al. (1998) found that when Upland cotton (*G. hirsutum*) cv. DPL-51 was grown in naturally lit plant growth chambers at 30/22°C day/night temperatures from sowing until flower bud production, and at 20/12, 25/17, 30/22, 35/27 and 40/32°C for 42 days after flower bud production, fruit retention was severely curtailed at the two higher temperatures compared with 30/22°C. Species/cultivars that retain fruits at high temperatures would be more productive both in the present-day cotton production environments and even more in future warmer world. Schrader et al. (2004) stated that high temperatures that plants are likely to experience inhibit photosynthesis.

Zhou et al. (2000) indicated that light duration is the key meteorological factor influencing the wheat-cotton cropping pattern and position of the bolls, while temperature had an important function on upper (node 7 to 9) and top (node 10) bolls, especially for double cropping patterns with early maturing varieties.

The objectives of this investigation were to study:

A- The effect of various climatic factors on the overall flower and boll production in Egyptian cotton. This could pave the way for formulating advanced predictions as for the effect of certain climatic conditions on cotton production of Egyptian cotton. It would be useful to minimize the deleterious effects of the factors through utilizing proper cultural practices which would limit and control their negative effects, and this will lead to an increase in cotton yield.

B- Further, this study aimed at predicting effects of climatic factors during different convenient intervals (in days) on cotton flower and boll production compared with daily observations. The study presents a rich effort focused on evaluating the efficacy of regression equations between cotton crop data and climatic data grouped at different time intervals, to determine the appropriate time scale for aggregating climate data to be used for predicting flower and boll production in cotton.

C- Furthermore, this study investigates and

collects information about the nature of the relationship between various climatic factors and cotton boll development and the 15-day period both prior to and after initiation of individual bolls of field grown cotton plants in Egypt. This could pave the way for formulating advanced predictions as for the effect of certain climatic conditions on production of Egyptian cotton. It would be useful to minimize the deleterious effects of the factors through utilizing proper cultural practices which would limit and control their negative effects, and this will lead to an improvement in cotton yield.

D- And provide information on the effect of various climatic factors and soil moisture status during the development stage on flower and boll production in Egyptian cotton. This could result in formulating advanced predictions as for the effect of certain climatic conditions on production of Egyptian cotton. Minimizing the deleterious effects of the factors through utilizing proper cultural practices will lead to improved cotton yield.

2. Data and Methods

Two uniform field trials were conducted at the experimental farm of the Agricultural Research Center, Ministry of Agriculture, Giza, Egypt (30°N, 31°: 28'E at an altitude of 19 m), using the cotton cultivar Giza 75 (*Gossypium barbadense* L.) in 2 successive seasons (I and II). The soil texture was a clay loam, with an alluvial substratum (pH = 8.07, 42.13% clay, 27.35% silt, 22.54% fine sand, 3.22% coarse sand, 2.94% calcium carbonate and 1.70% organic matter) (Sawan et al. 2010).

In Egypt, there are no rain-fed areas for cultivating cotton. Water for the field trials was applied using surface irrigation. Total water consumed during each of two growing seasons supplied by surface irrigation was about 6,000- $\text{m}^3 \text{ h}^{-1}$. The criteria used to determine amount of water applied to the crop depended on soil water status. Irrigation was applied when soil water content reached about 35% of field capacity (0-60 cm). In season I, the field was irrigated on 15 March (at planting), 8 April (first irrigation), 29 April, 17 May, 31 May, 14 June, 1 July, 16 July, and 12 August. In season II, the field was irrigated on 23 March (planting date), 20 April (first irrigation), 8 May, 22 May, 1 June, 18 June, 3 July, 20 July, 7 August and 28 August.

Techniques normally used for growing cotton in Egypt were followed. Each experimental plot contained 13 to 15 ridges to facilitate proper surface irrigation. Ridge width was 60 cm and length was 4 m. Seeds were sown on 15 and 23 March in seasons I and II, respectively, in hills 20 cm apart on one side of the ridge. Seedlings were thinned to 2 plants per hill 6 weeks after planting, resulting in a plant density of about 166,000 plants ha^{-1} . Phosphorus fertilizer was applied at a rate of 54 kg $\text{P}_2\text{O}_5 \text{ ha}^{-1}$ as calcium super phosphate during land preparation. Potassium fertilizer was applied at a rate of 57 kg $\text{K}_2\text{O} \text{ ha}^{-1}$ as potassium sulfate before the first irrigation (as a concentrated band close to the seed ridge). Nitrogen fertilizer was applied at a rate of 144 kg N ha^{-1} as ammonium nitrate in two equal doses: the first was applied after thinning just before the second irrigation and the second was applied before the third irrigation. Rates of phosphorus, potassium, and nitrogen fertilizer were the same in both seasons. These amounts were determined based on the use of soil tests (Sawan et al. 2010).

After thinning, 261 and 358 plants were randomly selected (precaution of border effect was taken into consideration by discarding the cotton plants in the first and last two hills of each ridge) from 9 and 11 inner ridges of the plot in seasons I, and II respectively. Pest control management was carried out on an-as-needed basis, according to the local practices performed at the experimental (Sawan et al. 2010).

Flowers on all selected plants were tagged in order to count and record the number of open flowers, and set bolls on a daily basis. The flowering season commenced on the date of the first flower appearance and continued until the end of flowering season (31 August). The period of whole September (30 days) until the 20th of October (harvest date) allowed a minimum of 50 days to develop mature bolls. In season I, the flowering period extended from 17 June to 31 August, whereas in season II, the flowering period was from 21 June to 31 August. Flowers produced after 31 August were not expected to form sound harvestable bolls, and therefore were not taken into account (Sawan et al. 2010).

For statistical analysis, the following data of the dependent variables were collected: number

of tagged flowers separately counted each day on all selected plants (Y_1), number of retained bolls obtained from the total daily tagged flowers on all selected plants at harvest (Y_2), and (Y_3) percentage of boll retention ([number of retained bolls obtained from the total number of daily tagged flowers in all selected plants at harvest]/[daily number of tagged flowers on each day in all selected plants] x 100).

As a rule, observations were recorded when the number of flowers on a given day was at least 5 flowers found in a population of 100 plants and this continued for at least five consecutive days. This rule omitted eight observations in the first season and ten observations in the second season. The number of observations (n) was 68 (23 June through 29 August) and 62 (29 June through 29 August) for the two seasons, respectively. Variables of the soil moisture status considered were, the day prior to irrigation, the day of irrigation, and the first and second days after the day of irrigation (Sawan et al. 2010).

The climatic factors (independent variables) considered were daily data of: maximum air temperature ($^{\circ}\text{C}$, X_1); minimum air temperature ($^{\circ}\text{C}$, X_2); maximum-minimum air temperature (diurnal temperature range) ($^{\circ}\text{C}$, X_3); evaporation (expressed as Piche evaporation) (mm day^{-1} , X_4); surface soil temperature, grass temperature or green cover temperature at 0600 h ($^{\circ}\text{C}$, X_5) and 1800 h ($^{\circ}\text{C}$, X_6); sunshine duration (h day^{-1} , X_7); maximum relative humidity (maxRH) (% , X_8), minimum relative humidity (minRH) (% , X_9) and wind speed (m s^{-1} , X_{10}) in season II only. The source of the climatic data was the Agricultural Meteorological Station of the Agricultural Research Station, Agricultural Research Center, Giza, Egypt. No rainfall occurred during the two growing seasons (Sawan et al. 2005).

Daily records of the climatic factors (independent variables), were taken for each day during production stage in any season including two additional periods of 15 days preceding and after the production stage (Sawan et al. 2005). Range and mean values of the climatic parameters recorded during the production stage for both seasons and overall data are listed in Table 1 (Sawan et al. 2006). Daily number of flowers and number of bolls per plant which survived till maturity (dependent variables) during the production stage in the two

seasons are graphically illustrated in Figures 1 and 2 (Sawan et al. 2005).

3. Results and Discussion

3.1. Response of flower and boll development to climatic factors on the anthesis day

Daily number of flowers and number of bolls per plant which survived to maturity (dependent variables) during the production stage of the two seasons (68 days and 62 days in the first and the second seasons, respectively) are graphically illustrated in Figures 1 and 2 (Sawan et al. 2005). The flower- and boll-curves reached their peaks during the middle two weeks of August, and then descended steadily till the end of the season. Specific differences in the shape of these curves in the two seasons may be due to the growth-reactions of environment, where climatic factors (Table 1) (Sawan et al. 2006) represent an important part of the environmental effects (Miller et al. 1996).

3.1.1. Correlation estimates:

Results of correlation coefficients [correlation and regression analyses were computed, according to Draper and Smith (1966) by means of the computer program SAS package (1985). between the initial group of independent variables and each of flower and boll production in the first and second seasons and the combined data of the two seasons are shown in Table 2 (Sawan et al. 2002).

The correlation values indicate clearly that evaporation is the most important climatic factor affecting flower and boll production as it showed the highest correlation value. This factor had a significant negative relationship with flower and boll production. Sunshine duration showed a significant negative relation with fruit production except for boll production in the first season, which was not significant. Maximum air temperature, temperature magnitude, and surface soil temperature at 1800 h, were also negatively correlated with flower and boll production in the second season and the combined data of the two seasons. Minimum humidity in the second season, the combined data of the two seasons, and maximum humidity in the first season were positively and highly correlated with flower and boll production. Minimum air temperature and

soil surface temperature at 0600 h showed low and insignificant correlation to flower and boll production (Sawan et al. 2002).

The negative relationship between evaporation with flower and boll production, means that high evaporation rate significantly reduces cotton flower and boll production. This may be due to greater plant water deficits when evaporation increases. Also, the negative relation between each of maximum temperature, temperature magnitude, surface soil temperature at 1800 h, or sunshine duration, with flower and boll production revealed that the increase in the values of these factors had a detrimental effect upon fruit production in Egyptian cotton. On the other hand, there was a positive correlation between each of maximum or minimum humidity with flower and boll production (Sawan et al. 2002).

Results obtained from the production stage of each season individually, and the combined data of the two seasons, indicate that relationships of some climatic variables with the dependent variables varied markedly from one season to another. This may be due to the differences between climatic factors in the two seasons as illustrated by the ranges and means shown in Table 1 (Sawan et al. 2006). For example, maximum temperature, minimum humidity and soil surface temperature at 1800 h did not show significant relations in the first season, while that trend differed in the second season. The effect of maximum humidity varied markedly from the first season to the second one. Where it was significantly correlated with the dependent variables in the first season, while the inverse pattern was true in the second season. This diverse effect may be due to the differences in the mean values of this factor in the two seasons; where it was, on average, about 86% in the first season, and about 72% on average in the second season, as shown in Table 1 (Sawan et al. 2006).

Boll retention ratio [(The number of retained bolls obtained from the total number of each daily tagged flowers in all selected plants at harvest/ Total number of daily tagged flowers of all selected plants) x 100] curves for both of the two seasons are shown in Figures 3 and 4 (Sawan et al. 2002). Also, these curves describe why the

shapes and patterns associated with the flower and boll curves for I and II seasons were different. It seems reasonable that the climatic data that were collected in these two experiments (I and II seasons) could provide adequate information for describing how these two seasons differed and how the crop responded accordingly (Sawan 2014 a & b).

These results indicate that evaporation is the most effective and consistent climatic factor affecting boll production. As the sign of the relationship was negative, this means that an increase in evaporation would cause a significant reduction in boll number. Thus, applying specific treatments such as an additional irrigation, and use of plant growth regulators, would decrease the deleterious effect of evaporation after boll formation and hence contribute to an increase in cotton boll production and retention, and the consequence is an increase in cotton yield (Sawan et al. 2002). In this connection, Moseley et al. (1994) stated that methanol has been reported to increase water use efficiency, growth and development of C_3 plants in arid conditions, under intense sunlight. In field trials cotton cv. DPL-50 (*Gossypium hirsutum*), was sprayed with a nutrient solution (1.33 lb N + 0.27 lb Fe + 0.27 lb Zn acre⁻¹) or 30% methanol solution at a rate of 20 gallons acre⁻¹, or sprayed with both the nutrient solution and methanol under two soil moisture regimes (irrigated and dry land). The foliar spray treatments were applied 6 times during the growing season beginning at first bloom. They found that irrigation (a total of 4.5 inches applied in July) increased lint yield across foliar spray treatments by 18%. Zhao and Oosterhuis (1997) reported that in a growth chamber when cotton (*Gossypium hirsutum* cv. Stoneville 506) plants were treated with the plant growth regulator PGR-IV (gibberellic acid, IBA and a proprietary fermentation broth) under water deficit stress and found significantly higher dry weights of roots and floral buds than the untreated water-stressed plants. They concluded that PGR-IV can partially alleviate the detrimental effects of water stress on photosynthesis and dry matter accumulation and improves the growth and nutrient absorption of growth chamber-grown cotton plants. Meek et al. (1999) in a field experiment in Arkansas found that application of 3 or 6 kg glycine betaine (PGR) ha⁻¹, to cotton plants had the potential for increasing yield in cotton exposed to mild water

stress.

3.1.2. Multiple linear regression equation:

By means of the multiple linear regression analysis, fitting predictive equations (having good fit) were computed for flower and boll production per plant using selected significant factors from the nine climatic variables studied in this investigation. Wind speed evaluated during the second season had no influence on the dependent variables. The equations obtained for each of the two dependent variables, i.e. number of flowers (Y_1) and bolls per plant (Y_2) in each season and for combined data from the two seasons (Table 2) (Sawan et al. 2002) are as follows:

First Season: (n = 68)

$$Y_1 = 21.691 - 1.968 X_4 - 0.241 X_7 + 0.216 X_8, R = 0.608^{**} \text{ and } R^2 = 0.3697,$$

While R^2 for all studied variables was 0.4022.

$$Y_2 = 15.434 - 1.633 X_4 + 0.159 X_8, R = 0.589^{**} \text{ and } R^2 = 0.3469 \text{ and } R^2 \text{ for all studied variables was } 0.3843.$$

Second Season: (n = 62)

$$Y_1 = 77.436 - 0.163 X_1 - 2.861 X_4 - 1.178 X_7 + 0.269 X_9, R = 0.644^{**}, R^2 = 0.4147.$$

$$Y_2 = 66.281 - 0.227 X_1 - 3.315 X_4 - 2.897 X_7 + 0.196 X_9, R = 0.629^{**}, R^2 = 0.3956.$$

In addition, R^2 for all studied variables was 0.4503 and 0.4287 for Y_1 and Y_2 equations respectively.

Combined data for the two seasons: (n = 130)

$$Y_1 = 68.143 - 0.827 X_4 - 1.190 X_6 - 2.718 X_7 + 0.512 X_9, R = 0.613^{**}, R^2 = 0.3758$$

$$Y_2 = 52.785 - 0.997 X_4 - 0.836 X_6 - 1.675 X_7 + 0.426 X_9, R = 0.569^{**}, R^2 = 0.3552$$

While R^2 for all studied variables was 0.4073 for Y_1 and 0.3790 for Y_2 .

Three climatic factors, i.e. minimum air temperature, surface soil temperature at 0600 h, and wind speed were not included in the equations since they had very little effect on production of cotton flowers and bolls. The sign of the partial regression coefficient for an independent

variable (climatic factor) indicates its effect on the production value of the dependent variable (flowers or bolls). This means that high rates of humidity and/or low values of evaporation will increase fruit production (Sawan et al. 2002).

3.1.3. Contribution of selected climatic factors to variations in the dependent variable:

Relative contributions (RC %) for each of the selected climatic factors to variation in flower and boll production is summarized in Table 3 (Sawan et al. 2002). Results in this table indicate that evaporation was the most important climatic factor affecting flower and boll production in Egyptian cotton. Sunshine duration is the second climatic factor of importance affecting production of flowers and bolls. Relative humidity and temperature at 1800 h were factors of lower contribution than evaporation and sunshine duration/day. Maximum temperature made a contribution less than the other affecting factors.

The highest contribution of evaporation to the variation in both flower and boll production (Sawan et al. 2002) can, however, be explained in the light of results found by Ward and Bunce (1986) in sunflower (*Helianthus annuus*). They stated that decreases of humidity at both leaf surfaces reduced photosynthetic rate of the whole leaf for plants grown under a moderate temperature and medium light level. Kaur and Singh (1992) found in cotton that flower number was decreased by water stress, particularly when applied at flowering. Seed cotton yield was about halved by water stress at flowering, slightly decreased by stress at boll formation, and not significantly affected by stress in the vegetative stage (6-7 weeks after sowing). Orgaz et al. (1992) in field experiments at Cordoba, SW Spain, grew cotton cultivars Acala SJ-C1, GC-510, Coker-310 and Jean cultivar at evapotranspiration (ET) levels ranging from 40 to 100% of maximum ET (ET_{max}) which were generated with sprinkler line irrigation. The water production function of Jean cultivar was linear; seed yield was 5.30 t ha⁻¹ at ET_{max} (820 mm). In contrast, the production function of the three other cultivars was linear up to 85% of ET_{max} , but leveled off as ET approached ET_{max} (830 mm) because a fraction of the set bolls did not open by harvest at high ET levels. These authors concluded that it is possible to

define an optimum ET deficit for cotton based on cultivar earliness, growing-season length, and availability of irrigation water.

The negative relationship between sunshine duration and cotton production may be due to the fact that the species of *Gossypium* used is known to be a short day plant (Hearn and Constable 1984), so, an increase of sunshine duration above that needed for cotton plant growth will decrease flower and boll production. Oosterhuis (1997) studied the reasons for low and variable cotton yields in Arkansas, with unusually high insect pressures and the development of the boll load during an exceptionally hot and dry August. Solutions to the problems are suggested i.e. selection of tolerant cultivars, effective and timely insect and weed control, adequate irrigation regime, use of proper crop monitoring techniques and application of plant growth regulators.

3.2. Appropriate time scale for aggregating climatic data to predict flowering and boll setting behavior of cotton

Statistical Analysis

Statistical analysis was conducted using the procedures outlined in the general linear model (GLM, SAS Institute, Inc. 1985). Data of dependent and independent variables, collected for each day of the production stage (60 days in each season), were summed up into intervals of 2, 3, 4, 5, 6 or 10 days. Data from these intervals were used to compute relationships between the dependent variables (flower and boll setting and boll retention) and the independent variables (climatic factors) in the form of simple correlation coefficients for each season. Comparisons between the values of "r" were done to determine the best interval of days for determining effective relationships. The α -level for significance was $P \leq 0.15$. The climatic factors attaining a probability level of significance not exceeding 0.15 were deemed important (affecting the dependent variables), selected and combined with dependent variable in multiple regression analysis to obtain a convenient predictive equation (Cady and Allen 1972). Multiple linear regression equations (using stepwise method) comprising selected predictive variables were computed for the determined interval and coefficients of multiple determinations (R^2) were calculated to measure the efficiency of the

regression models in explaining the variation in data. Correlation and regression analyses were computed according to Draper and Smith (1966) (Sawan et al. 2006).

3.2.1. Correlation estimates

Significant simple correlation coefficients were estimated between the production variables and studied climatic factors for different intervals of days (combined data of the 2 seasons) (Table 4) (Sawan et al. 2006).

Evaporation was the most important climatic factor affecting flower and boll production in Egyptian cotton. The negative correlation means that high evaporation ratio significantly reduced flower and boll production. High evaporation rates could result in water stress that would slow growth and increase shedding rate of flowers and bolls (Sawan et al. 2006). Kaur and Singh (1992) found in cotton that flower number was decreased by water stress, particularly when existing at flowering stage. Seed cotton yield was decreased by about 50% when water stress was present at flowering stage, slightly decreased by stress at boll formation stage, and not significantly affected by stress in the vegetative stage (6-7 weeks after sowing).

The second most important climatic factor was minimum humidity, which had a high positive correlation with flower and boll production, and retention ratio. The positive correlation means that increased humidity would bring about better boll production.

The third most important climatic factor in our study was sunshine duration, which showed a significant negative relationship with flower and boll production only. The negative relationship between sunshine duration and cotton production may be due to the fact that the species of the genus *Gossypium* are known to be short day plants (Hearn and Constable 1984), so, an increase of sunshine duration above that sufficient to attain good plant growth will decrease flower and boll production. Bhatt (1977) found that exposure to daylight over 14 hours and high day temperature, individually or in combination, delayed flowering of the Upland cotton cv. J34. Although average sunshine duration in our study was only 11.7 h, yet it could reach 13 h, which, in combination with

high maximum temperatures (up to 38.8°C), may have adversely affected reproductive growth.

Maximum air temperature, temperature magnitude and surface soil temperature at 1800 h show significant negative relationships with flower and boll production only. Meanwhile, the least important factors were surface soil temperature at 0600 h and minimum air temperature. Our results indicate that evaporation was the most effective climatic factor affecting cotton boll production. As the sign of the relationship was negative, this means that an increase in evaporation caused a significant reduction in boll number (Sawan et al. 2006). Thus, applying specific treatments, such as an additional irrigation or the use of plant growth regulators (PGR) that would decrease the deleterious effect of evaporation after boll formation, could contribute to an increase in cotton boll production and retention, and consequently an increase in cotton yield. In this connection, Meek et al. (1999) in a field experiment in Arkansas found that application of 3 or 6 kg glycine betaine (PGR) ha⁻¹ to cotton plants under mild water stress increased yield.

Comparing results for the different intervals of days with those from daily observation (Table 4) (Sawan et al. 2006), the 5-day interval appeared to be the most suitable interval, which actually revealed a more solid and more obvious relationships between climatic factors and production characters. This was in fact indicated by the higher R² values obtained when using the 5-day intervals. The 5-day interval may be the most suitable interval for diminishing the daily fluctuations between the factors under study to clear these relations comparing with the other intervals. However, it seems that this conception is true provided that the fluctuations in climatic conditions are limited or minimal. Therefore, it would be the most efficient interval used to help circumvent the unfavorable effect of climatic factors. This finding gives researchers and producers a chance to deal with condensed rather than daily weather data.

3.2.2. Regression models

Multiple linear regression equations were estimated using the stepwise multiple regression technique to express the relation between cotton production variables [number of flowers (Y_1);

bolls per plant (Y_2); and boll retention ratio (Y_3)] and the studied climatic factors (Table 5) (Sawan et al. 2006).

Evaporation and surface soil temperature at 1800 h, sunshine duration and minimum humidity accounted for a highly significant amount of variation ($P < 0.05$) in cotton production variables, with the equation obtained for the 5-day interval showing a high degree of certainty. The R² values for the 5-day interval were higher than those obtained from daily data for each of the cotton production variables. Also, the 5-day interval gave more efficient and stable estimates than the other studied intervals (data not shown) (Sawan et al. 2006). The R² values for these equations clearly indicate the importance of such equations since the climatic factors involved explained about 59 to 62% of the variation found in the dependent variables.

During the production stage, an accurate weather forecast for the next 10 days would provide an opportunity to avoid any adverse effect for weather factors on cotton production through applying appropriate cultural practices such as adequate irrigation regime or utilization of plant growth regulators. This proposal would be true if the fluctuations in weather conditions were not extreme. Our recommendation would be the accumulation 5-day climatic data, and use this information to select the adequate cultural practices (such as an additional irrigation or utilization of plant growth regulators) that would help circumvent the unfavorable effects of climatic factors. In case of sharp fluctuations in climatic factors, data could be collected daily, and when stability of climatic conditions is restored, the 5-day accumulation of weather data could be used again (Sawan et al. 2006).

3.3. Response of flower and boll development to climate factors before and after anthesis day

The effects of specific climatic factors during both pre- and post-anthesis periods on boll production and retention are mostly unknown. However, by determining the relationship of climatic factors with flower and boll production and retention, the overall level of production can be possibly predicted. Thus, an understanding of these relationships may help physiologists to determine control mechanisms of production in

cotton plants (Sawan et al. 2005). Daily records of the climatic factors (independent variables), were taken for each day during production stage in any season including two additional periods of 15 days before and after the production stage (Table 6) (Sawan et al. 2005).

In each season, the data of the dependent and independent variables (68 and 62 days) were regarded as the original file (a file which contains the daily recorded data for any variable during a specific period). Fifteen other files before and another 15 after the production stage were obtained by fixing the dependent variable data, while moving the independent variable data at steps each of 1 day (either before or after production stage) in a manner similar to a sliding role (Sawan et al. 2005). The following is an example (in the first season):

Thus, the climate data were organized into records according to the complete production stage (68 days the first year and 62 days the second year) and 15 day, 14 day, 13 day,....and 1 day periods both before and after the production stage. This produced 31 climate periods per year that were analyzed for their relationships with cotton flowering and boll production (Sawan et al. 2005).

3.3.1. Correlation estimates

A. Results of the correlation between climatic factors and each of flower and boll production during the 15 day periods before flowering day (Tables 7 and 8) revealed the following (Sawan et al. 2005):

First season

File	Data of any dependent variable (for each flowers and bolls)		Any independent variable (for each climatic factors)			
	Production stage		In case of original file and files before production stage		In case of original file and files after production stage	
	Date	Days	Date	Days	Date	Days
Original file	23 Jun-29 Aug	68	23 Jun-29 Aug	68	23 Jun-29 Aug	68
1 st new file	23 Jun-29 Aug	68	22 Jun-28 Aug	68	24 Jun-30 Aug	68
2 nd new file	23 Jun-29 Aug	68	21 Jun-27 Aug	68	25 Jun-31 Aug	68
15 th new file	23 Jun-29 Aug	68	8 Jun-14 Aug	68	8 Jul -13 Sept	68

Daily evaporation and sunshine duration showed consistent negative and statistically significant correlations with both flower and boll production for each of the 15 moving window periods before anthesis (Table 7). Evaporation appeared to be the most important climate factor affecting flower and boll production.

Daily maximum and minimum humidity showed consistent positive and statistically significant correlations with both flower and boll production in most of the 15 moving window periods before anthesis (Table 7) (Sawan et al. 2005). Maximum daily temperature showed low but significant negative correlation with flower production during the 2-5, 8, and 10 day periods before anthesis. Minimum daily temperatures generally showed insignificant correlation with both production variables. The diurnal temperature range showed few correlations with flower and boll production. Daily soil surface temperature at 0600 h showed a significant positive correlation with boll production during the period extending from the 11-15 day period before anthesis, while its effect on flowering was confined only to the 12 and the 15 day periods prior anthesis. Daily soil surface temperature at 1800 h showed a significant negative correlation with flower production during the 2-10 day periods before anthesis.

Second season

Daily Evaporation, the diurnal temperature range, and sunshine duration were negatively and significantly correlated with both flower and boll production in all the 15 day periods, while maximum daily temperature was negatively and

significantly related to flower and boll formation during the 2- 5 day periods before anthesis (Table 8) (Sawan et al. 2005).

Minimum daily temperature showed positive and statistically significant correlations with both production variables only during the 9-15 day periods before anthesis, while daily minimum humidity showed the same correlation trend in all the 15 moving window periods before anthesis. Daily soil surface temperature at 0600 h was positively and significantly correlated with flower and boll production for the 12, 14, and 15 day periods prior to anthesis only. Daily soil surface temperature at 1800 h showed negative and significant correlations with both production variables only during the first and second day periods before flowering. Daily maximum humidity showed insignificant correlation with both flower and boll production except for one day period only (the 15th day). Generally, the results in the two seasons indicated that daily evaporation, sunshine duration and minimum humidity were the most effective and consistent climatic factors, which exhibited significant relationships with the production variables for all the 15 day periods before anthesis in both seasons (Sawan et al. 2005).

The factors in this study which had been found to be associated with boll development are the climatic factors that would influence water loss between plant and atmosphere (low evaporation demand, high humidity, and shorter solar duration). This can lead to direct effects on the fruiting forms themselves and inhibitory effects on mid-afternoon photosynthetic rates even under well-watered conditions. Boyer et al. (1980) found that soybean plants with ample water supplies can experience water deficits due to high transpiration rates. Also, Human et al. (1990) stated that, when sunflower plants were grown under controlled temperature regimes, water stress during budding, anthesis and seed filling, the CO₂ uptake rate per unit leaf area as well as total uptake rate per plant, significantly diminished with stress, while this effect resulted in a significant decrease in yield per plant.

B. The correlation between climatic factors and each of boll production and boll retention over a period of 15 day periods after flowering (boll setting) day (Tables 9 and 10) (Sawan et

al. 2005) revealed the following:

First season

Daily evaporation showed significant negative correlation with number of bolls for all the 15 day periods after flowering (Table 9). Meanwhile its relationship with retention ratio was positive and significant in the 9-15 day periods after flowering. Daily sunshine duration was positively and significantly correlated with boll retention ratio during the 5-13 day periods after flowering. Daily maximum humidity had a significant positive correlation with the number of bolls during the first 8 day periods after flowering, while daily minimum humidity had the same correlation for only the 11, and 12 day periods after flowering. Daily maximum and minimum temperatures and the diurnal temperature range, as well as soil surface temperature at 1800 did not show significant relationships with both number of bolls and retention ratio. Daily soil surface temperature at 0600 h had a significant negative correlation with boll retention ratio during the 3-7 day periods after anthesis.

Second season

Daily evaporation, soil surface temperature at 1800 h, and sunshine duration had a significant negative correlation with number of bolls in all the 15 day periods after anthesis (Table 10) (Sawan et al. 2005). Daily maximum and minimum temperatures and the diurnal temperature range, and soil surface temperature at 0600 h had a negative correlation with boll production. Their significant effects were observed during the 1, and 10-15 day periods for maximum temperature, and the 1-5, and 9-12 day periods for the diurnal temperatures range. Meanwhile, the daily minimum temperature and soil surface temperature at 0600 h had a significant negative correlation only during the 13-15 day periods. Daily minimum humidity had a significant positive correlation with number of bolls during the first 5 day periods, and the 9-15 day periods after anthesis. Daily maximum humidity showed no significant relation to number of bolls produced, and further no significant relation was observed between any of the studied climatic factors and boll retention ratio.

The results in the two seasons indicated that evaporation and humidity, followed by

sunshine duration had obvious correlation with boll production. From the results obtained, it appeared that the effects of air temperature, and soil surface temperature tended to be masked in the first season, i.e. did not show any significant effects in the first season on the number of bolls per plant. However, these effects were found to be significant in the second season. These seasonal differences in the impacts of the previously mentioned climatic factors on the number of bolls per plant are most likely ascribed to the sensible variation in evaporation values in the two studied seasons where their means were 10.2 mm.d^{-1} and 5.9 mm.d^{-1} in the first and second seasons, respectively (Sawan et al. 2005).

There is an important question here concerning, if there is a way for forecasting when evaporation values would mask the effect of the previous climatic factors. The answer would be possibly achieved through relating humidity values to evaporation values which are naturally liable to some fluctuations from one season to another (Sawan et al. 2005). It was found that the ratio between the mean of maximum humidity and the mean of evaporation in the first season was $85.8/10.2 = 8.37$, while in the second season this ratio was 12.4. On the other hand, the ratio between the mean minimum humidity and the mean of evaporation in the first season was $30.8/10.2 = 3.02$, while in the second season this ratio was 6.75 (Table 9) (Sawan et al. 2005). From these ratios it seems that minimum humidity which is closely related to evaporation is more sensitive than the ratio between maximum humidity and evaporation. It can be seen from the results and formulas that when the ratio between minimum humidity and evaporation is small (3:1), the effects of air temperature, and soil surface temperature were hindered by the effect of evaporation, i.e. the effect of these climatic factors were not significant. However, when this ratio is high (6:1), the effects of these factors were found to be significant. Accordingly, it could be generally stated that the effects of air, and soil surface temperatures could be masked by evaporation when the ratio between minimum humidity and evaporation is less than 4:1 (Sawan et al. 2005).

Evaporation appeared to be the most important climatic factor (in each of the 15-day periods both prior to and after initiation of individual

bolls) affecting number of flowers or harvested bolls in Egyptian cotton. High daily evaporation rates could result in water stress that would slow growth and increase shedding rate of flowers and bolls. The second most important climatic factor in our study was humidity. Effect of maximum humidity varied markedly from the first season to the second one, where it was significantly correlated with the dependent variables in the first season, while the inverse pattern was true in the second season. This diverse effect may be due to the differences in the values of this factor in the two seasons; where it was on average 87% in the first season, and only 73% in the second season (Table 6) (Sawan et al. 2005). Also, was found that, when the average value of minimum humidity exceeded the half average value of maximum humidity, the minimum humidity can substitute the maximum humidity on affecting number of flowers or harvested bolls. In the first season (Table 10) the average value of minimum humidity was less than half of the value of maximum humidity ($30.2/85.6 = 0.35$), while in the second season it was higher than half of maximum humidity ($39.1/72.9 = 0.54$).

The third most important climatic factor in our study was sunshine duration, which showed a significant negative relationship with boll production. The r values of (Tables 7-10) (Sawan et al. 2005) indicated that the relationship between the dependent and independent variables preceding flowering (production stage) generally exceeded in value the relationship between them during the entire and late periods of production stage. In fact, understanding the effects of climatic factors on cotton production during the previously mentioned periods would have marked consequences on the overall level of cotton production, which could be predictable depending on those relationships.

3.3.2. Regression models

An attempt was carried out to investigate the effect of climatic factors on cotton production via prediction equations including the important climatic factors responsible for the majority of total variability in cotton flower and boll production. Hence, regression models were established using the stepwise multiple regression technique to express the relationship between each of the number of flowers and bolls/plant and boll

retention ratio (Y), with the climatic factors, for each of the a) 5, b) 10, and c) 15 day periods either prior to or after initiation of individual bolls (Tables 11 and 12) (Sawan et al. 2005).

Concerning the effect of prior days the results indicated that evaporation, sunshine duration, and the diurnal temperature range were the most effective and consistent climatic factors affecting cotton flower and boll production (Table 11). The fourth effective climatic factor in this respect was minimum humidity. On the other hand, for the periods after flower the results obtained from the equations (Table 12) indicated that evaporation was the most effective and consistent climatic factor affecting number of harvested bolls.

Regression models obtained demonstrate of each independent variable under study as an efficient and important factor (Sawan et al. 2005). Meanwhile, they explained a sensible proportion of the variation in flower and boll production, as indicated by their R^2 , which ranged between 0.14-0.62, where most of R^2 prior to flower opening were about 0.50 and after flowering all but one are less than 0.50. These results agree with Miller et al. (1996) in their regression study of the relation of yield with rainfall and temperature. They suggested that the other 0.50 of variation related to management practices, which can be the same in this study. Also, the regression models indicated that the relationships between the number of flowers and bolls per plant and the studied climatic factors for the 15 day period before or after flowering (Y_3) in each season explained the highly significant magnitude of variation ($P < 0.05$). The R^2 values for the 15 day periods before and after flowering were higher than most of those obtained for each of the 5 and the 10 day periods before or after flowering. This clarifies that the effects of the climatic factors during the 15 day periods before or after flowering are very important for Egyptian cotton boll production and retention. Thus, an accurate climatic forecast for the effect of these 15 day periods provides an opportunity to avoid any possible adverse effects of unusual climatic conditions before flowering or after boll formation by utilizing additional treatments and/or adopting proper precautions to avoid flower and boll reduction.

The main climatic factors from this study (Sawan

et al. 2005) affecting the number of flowers and bolls, and by implication yield, is evaporation, sunshine duration and minimum humidity, with evaporation (water stress) being by far the most important factor. Various activities have been suggested to partially overcome water stress. Temperature conditions during the reproduction growth stage of cotton in Egypt do not appear to limit growth even though they are above the optimum for cotton growth (Sawan 2013). This is contradictory to the finding of Holaday et al. (1997). A possible reason for that contradiction is that the effects of evaporation rate and humidity were not taken into consideration in the research studies conducted by other researchers in other countries. The matter of fact is that temperature and evaporation are closely related to each other to such an extent that the higher evaporation rate could possible mask the effect of temperature (Sawan 2014a). Sunshine duration and minimum humidity appeared to have secondary effects, yet they are in fact important players. The importance of sunshine duration has been alluded to by Moseley et al. (1994) and Oosterhuis (1997). Also, Mergeai and Demol (1991) found that cotton yield was assisted by intermediate relative humidity.

3.4. Cotton (*Gossypium barbadense*) flower and boll production as affected by climatic factors and soil moisture status

Basic Variables

A. Dependant variables as defined above: (Y_1) and (Y_2) (Sawan et al. 2010).

B. Independent variables (X_s):

1. Irrigation on day 1 = 1. Otherwise, enter 0.0 (soil moisture status) (X_1)
2. The first and second days after the day of irrigation (soil moisture status) = 1. Otherwise, enter 0.0 (X_2).
3. The day prior to the day of irrigation (soil moisture status) to check for possible moisture deficiency on that day = 1. Otherwise, enter 0.0 (X_3).
4. Number of days during days 1 (day of flowering)-12 (after flowering) that temperature equaled or exceeded 37.5 °C (high temperature)

(X4).

5. Range of temperature (diurnal temperature) [°C] on day 1 (day of flowering) (X5).

6. Broadest range of temperature [°C] over days 1 (day of flowering)-12 (after flowering) (X6).

7. Minimum relative humidity (minRH) [%] during day 1 (day of flowering) (X7).

8. Maximum relative humidity (maxRH) [%] during day 1 (day of flowering) (X8).

9. Minimum relative humidity (minRH) [%] during day 2 (after flowering) (X9).

10. Maximum relative humidity (maxRH) [%] during day 2 (after flowering) (X10).

11. Largest maximum relative humidity (maxRH) [%] on days 3-6 (after flowering) (X11).

12. Lowest minimum relative humidity (minRH) [%] on days 3-6 (after flowering) (X12).

13. Largest maximum relative humidity (maxRH) [%] on days 7-12 (after flowering) (X13).

14. Lowest minimum relative humidity (minRH) [%] on days 7-12 (after flowering) (X14).

15. Lowest minimum relative humidity (minRH) [%] on days 50-52 (after flowering) (X15).

16. Daily light period (hour) (X16).

Statistical analysis

Simple correlation coefficients between the initial group of independent variables (climatic factors and soil moisture status) (X's) and the corresponding dependent variables (Y's) were computed for each season and the combined data of the two seasons. These correlation coefficients helped determine the significant climatic factors and soil moisture status affecting the cotton production variables. The level for significance was $P \leq 0.15$. Those climatic factors and soil moisture status attaining a probability level of significance not exceeding 0.15 were deemed important (affecting the dependent variables) (Sawan et al. 2010). Those factors were combined with dependent variables in multiple regression analysis to obtain a predictive model as described by Cady and Allen (1972).

Multiple linear regression equations (using the stepwise method) comprising selected predictive variables were computed for the determined interval. Coefficients of multiple determinations (R^2) were calculated to measure the efficiency of the regression models in explaining the variation in data. Correlation and regression analysis were computed according to Draper and Smith (1985) using the procedures outlined in the general linear model (GLM) (SAS Institute 1985).

3.4.1. Correlation estimates

Simple correlation coefficients between the independent variables and the dependent variables for flower and boll production in each season and combined data of the two seasons are shown in Tables 13-15 (Sawan et al. 2010). The simple correlation values indicated clearly that relative humidity was the most important climatic factor. Relative humidity also had a significant positive relationship with flower and boll production; except for lowest minRH on days 50-52 (after flowering). Flower and boll production were positively and highly correlated with the variables of largest maxRH (X11, X13) and lowest minRH (X14, X15) in the first season, minRH (X7, X9), largest maxRH (X11), and lowest minRH (X12, X14, X15) in the second season, and the combined data of the two seasons. Effect of maxRH varied markedly from the first to the second season. MaxRH was significantly correlated with the dependent variables in the first season, while the inverse pattern was true in the second season. This diverse effect may be best explained by the differences of 87% in the first season, and only 73% in the second season (Table 1). Also, when the average value of minRH exceeded the half average value of maxRH, the minRH can substitute for the maxRH on affecting number of flowers or harvested bolls. In the first season (Table 1) the average value of minRH was less than half of the value of maxRH ($30.2/85.6 = 0.35$), while in the second season it was higher than half of maxRH ($39.1/72.9 = 0.54$). Sunshine duration (X16) showed a significant negative relation with fruit production in the first and second seasons and the combined data of the two seasons except for boll production in the first season, which was not significant. Flower and boll production were negatively correlated in the second season and the combined data of the two seasons with the number of days

during days 1 -12 that temperature equaled or exceeded 37.5 °C (X4), range of temperature (diurnal temperature) on flowering day (X5) and broadest range of temperature over days 1-12 (X6). The soil moisture status showed low and insignificant correlation with flower and boll production. The positive relationship between relative humidity with flower and boll production means that low relative humidity rate reduces significantly cotton flower and boll production. This may be due to greater plant water deficits when relative humidity decreases. Also, the negative relationship between the variables of maximum temperature exceeding 37.5 °C (X4), range of diurnal temperature on flowering (X5), and sunshine duration (X16) with flower and boll production revealed that the increased values of these factors had a detrimental effect upon Egyptian cotton fruit production. Results obtained from the production stage of each season, and the combined data of the two seasons showed marked variability in the relationships of some climatic variables with the dependent variables. This may be best explained by the differences between climatic factors in the two seasons as illustrated by the ranges and means shown in Table 1. For example, maximum temperature exceeding 37.5 °C (X4) and minRH did not show significant relations in the first season, while that trend differed in the second season. These results indicated that relative humidity was the most effective and consistent climatic factor affecting boll production. The second most important climatic factor in our study was sunshine duration, which showed a significant negative relationship with boll production.

3.4.2. Multiple linear regression models, beside contribution of climatic factors and soil moisture status to variations in the dependent variables

Regression models were established using the stepwise multiple regression technique to express the relationship between the number of flowers and bolls per plant⁻¹ (Y) with the climatic factors and soil moisture status (Table 16). Relative humidity (%) was the most important climatic factor affecting flower and boll production in Egyptian cotton [minRH during day 1 (X7), minRH during day 2 (X9), largest maxRH on days 3-6 (X11), lowest minRH on days 3-6 (X12), largest maxRH on days 7-12

(X13), lowest minRH on days 7-12 (X14) and lowest minRH on days 50-52 (X15)]. Sunshine duration (X16) was the second climatic factor of importance affecting production of flowers and bolls. Maximum temperature (X4), broadest range of temperature (X6) and soil moisture status (X1) made a contribution affecting flower and boll production. The soil moisture variables (X2, X3), and climatic factors (X5, X8, X10) were not included in the equations since they had very little effects on production of cotton flowers and bolls.

Relative humidity showed the highest contribution to the variation in both flower and boll production (Table 16). This finding can be explained in the light of results found by Ward and Bunce (1986) in sunflower (*Helianthus annuus*). They stated that decreases of relative humidity on both leaf surfaces reduced photosynthetic rate of the whole leaf for plants grown under a moderate temperature and medium light level.

Reddy et al. (1993) found that cotton (*Gossypium hirsutum*) fruit retention decreased rapidly as the time of exposure to 40°C increased. Gutiérrez and López (2003) studied the effects of heat on the yield of cotton in Andalusia, Spain, during 1991-98, and found that high temperatures were implicated in the reduction of unit production. There was a significant negative relationship between average production and number of days with temperatures greater than 40°C and the number of days with minimum temperatures greater than 20°C. Wise et al. (2004) indicated that restrictions to photosynthesis could limit plant growth at high temperature in a variety of ways. In addition to increasing photorespiration, high temperatures (35-42°C) can cause direct injury to the photosynthetic apparatus. Both carbon metabolism and thylakoid reactions have been suggested as the primary site of injury at these temperatures.

Regression models obtained explained a sensible proportion of the variation in flower and boll production, as indicated by their R², which ranged between 0.53-0.72. These results agree with Miller et al. (1996) in their regression study of the relation of yield with rainfall and temperature. They suggested that the other R² 0.50 of variation was related to management practices, which coincide with the findings of this study. Thus,

an accurate climatic forecast for the effect of the 5-7 day period during flowering may provide an opportunity to avoid possible adverse effects of unusual climatic conditions before flowering or after boll formation by utilizing additional treatments and/or adopting proper precautions to avoid flower and boll reduction (Sawan 2013).

4. Conclusions

Evaporation, sunshine duration, relative humidity, surface soil temperature at 1800 h, and maximum temperature, were the most significant climatic factors affecting flower and boll production of Egyptian cotton. Also, it could be concluded that during the 15-day periods both prior to and after initiation of individual bolls, evaporation, minimum relative humidity and sunshine duration, were the most significant climatic factors affecting cotton flower and boll production and retention in Egyptian cotton. The negative correlation between each of evaporation and sunshine duration with flower and boll formation along with the positive correlation between minimum relative humidity value and flower and boll production, indicate that low evaporation rate, short period of sunshine duration and high value of minimum humidity would enhance flower and boll formation. The 5-day interval was found to give adequate and sensible relationships between climatic factors and cotton production growth under Egyptian conditions when compared with other intervals and daily observations. It may be concluded that the 5-day accumulation of climatic data during the production stage, in the absence of sharp fluctuations in these factors, could be satisfactorily used to forecast adverse effects on cotton production and the application of appropriate production practices circumvent possible production shortage.

Finally, the early prediction of possible adverse effects of climatic factors might modify their effect on production of Egyptian cotton. Minimizing deleterious effects through the application of proper management practices, such as, adequate irrigation regime, and utilization of specific plant growth regulators could limit the negative effects of some climatic factors.

References

- Barbour MM, Farquhar GD (2000) Relative humidity- and ABA-induced variation in carbon and oxygen isotope ratios of cotton leaves. *Plant, Cell and Environment* 23: 473-485.
- Bhatt JG (1977) Growth and flowering of cotton (*Gossypium hirsutum* L.) as affected by daylength and temperature. *Journal of Agricultural Science* 89: 583-588.
- Boyer JS, Johnson RR, Saupe SG (1980) Afternoon water deficits and grain yields in old and new soybean cultivars. *Agron J* 72: 981-986.
- Burke JJ, Mahan JR, Hatfield JL (1988) Crop specific thermal kinetic windows in relation to wheat and cotton biomass production. *Agron J* 80: 553-556.
- Cady FB, Allen DM (1972) Combining experiments to predict future yield data. *Agron J* 64: 211-214.
- Draper NR, Smith H (1966) *Applied Regression Analysis*. John Wiley & Sons Ltd., New York, NY. 407 pp.
- El-Zik KM (1980) The cotton plant - its growth and development. *Western Cotton Prod. Conf. Summary Proc.*, Fresno, CA, p. 18-21.
- Guinn G (1982) Causes of square and boll shedding in cotton. *USDA Tech. Bull.* 1672. USDA, Washington, DC.
- Gutiérrez Mas JC, López M (2003) Heat, limitation of yields of cotton in Andalucía. *Agricultura, Revista Agropecuaria* 72: 690-692.
- Hearn AB, Constable GA (1984) *The Physiology of Tropical Food Crops*. Chapter 14: Cotton. P. 495-527 (Edited by Goldsworth, P.R.; Fisher, N.M.), John Wiley & Sons Ltd., NY. 664 pp.
- Hodges HF, Reddy KR, McKinion JM, Reddy VR (1993) Temperature effects on cotton. *Bulletin Mississippi Agricultural and Forestry Experiment Station No.* 990: 15.
- Holaday AS, Haigler CH, Srinivas NG, Martin LK, Taylor JG (1997) Alterations of leaf photosynthesis and fiber cellulose synthesis by cool night temperatures. In *Proceedings Beltwide Cotton Conferences*, January 6-10, New Orleans, LA, National Cotton Council, TN pp 1435-1436.
- Human JJ, Du Toit D, Bezuidenhout HD, De Bruyn LP (1990) The influence of plant water stress on net photosynthesis and yield of sunflower (*Helianthus annuus* L.). *J Agron Crop Sci* 164: 231-241.
- Kaur R, Singh OS (1992) Response of growth stages of cotton varieties to moisture stress. *Indian J Plant Physiol* 35: 182-185.
- Meek CR, Oosterhuis DM, Steger AT (1999) Drought tolerance and foliar sprays of glycine betaine. In *Proceedings Beltwide Cotton Conferences*, January 3-7, Orlando, FL, USA. Memphis, USA. National Cotton Council, 559-561.
- Mergeai G, Demol J (1991) Contribution to the study

- of the effect of various meteorological factors on production and quality of cotton (*Gossypium hirsutum* L.) fibers. *Bulletin des Recherches Agronomiques de Gembloux* 26: 113-124.
- Miller JK, Krieg DR, Paterson RE (1996) Relationship between dryland cotton yields and weather parameters on the Southern Hig Plains. In *Proceedings Beltwide Cotton Conferences*, January 9-12, Nashville, TN, USA, Memphis, USA, National Cotton Council, 1165-1166.
- Moseley D, Landivar JA, Locke D (1994) Evaluation of the effect of methanol on cotton growth and yield under dry-land and irrigated conditions. In *Proceedings Beltwide Cotton Conferences*, January 5-8, San Diego, CA, USA. Memphis, USA. National Cotton Council, 1293-1294.
- Oosterhuis DM (1997) Effect of temperature extremes on cotton yields in Arkansas. In *Proceedings of the Cotton Research Meeting*, held at Monticello, Arkansas, USA, February 13 [Edited by Oosterhuis, D.M.; Stewart, J.M.]. Special Report-Agricultural Experiment Station, Division of Agriculture, University of Arkansas, No. 183: 94-98.
- Orgaz F, Mateos L, Fereres E (1992) Season length and cultivar determine the optimum evapotranspiration deficit in cotton. *Agron J* 84: 700-706.
- Reddy KR, Hodges HF, McKinion JM (1993) Temperature effects on Pima cotton leaf growth. *Agron J* 85: 681-686.
- Reddy KR, Robana RR, Hodges HF, Liu XJ, McKinion JM (1998) Interactions of CO₂ enrichment and temperature on cotton growth and leaf characteristics. *Environ Exp Bot* 39: 117-129.
- SAS Institute, Inc (1985) *SAS User's Guide: Statistics*. 5th ed. SAS Institute, Inc., Cary, NC. pp. 433-506.
- Sawan ZM (2013) Studying the relationship between climatic factors and cotton production by different applied methods. *Journal of Stress Physiology & Biochemistry* 9: 251-278.
- Sawan ZM (2014a) Climatic factors: evaporation, sunshine, relative humidity, soil and air temperature and cotton production. *Annual Research & Review in Biology*, 4: 2835-2855.
- Sawan ZM (2014b) Nature relation between climatic variables and cotton production. *Journal of Stress Physiology & Biochemistry* 9: 251-278.
- Sawan ZM, Hanna LI, Gad El Karim GhA, McCuistions WL (2002) Relationships between climatic factors and flower and boll production in Egyptian cotton (*Gossypium barbadense*). *Journal of Arid Environment* 52: 499-516.
- Sawan ZM, Hanna LI, McCuistions WL (2005) Response of flower and boll development to climatic factors before and after anthesis in Egyptian cotton. *Climate Research* 29: 167-179.
- Sawan ZM, Hanna LI, McCuistions WL (2006) Appropriate time scale for aggregating climatic data to predict flowering and boll setting behaviour of cotton in Egypt. *Communication in Biometry and Crop Science* 1: 11-19.
- Sawan ZM, Hanna LI, McCuistions WL, Foote RJ (2010) Egyptian cotton (*Gossypium barbadense*) flower and boll production as affected by climatic factors and soil moisture status. *Theoretical and Applied Climatology* 99: 217-227.
- Schrader SM, Wise RR, Wacholtz WF, Ort DR, Sharkey TD (2004) Thylakoid membrane responses to moderately high leaf temperature in Pima Cotton. *Plant, Cell and Environment* 27: 725-735.
- Ward DA, Bunce JA (1986) Responses of net photosynthesis and conductance to independent changes in the humidity environments of the upper and lower surfaces of leaves of sunflower and soybean. *J Exper Botany* 37: 1842-1853.
- Wise RR, Olson AJ, Schrader SM, Sharkey TD (2004) Electron transport is the functional limitation of photosynthesis in field-grown Pima cotton plants at high temperature. *Plant, Cell and Environment* 27: 717-724.
- Xiao J-F, Liu Z-G, Yu X-G, Zhang J-Y, Duan A-W (2000) Effects of different water application on lint yield and fiber quality of cotton under drip irrigation. *Acta Gossypii Sinica* 12: 194-197.
- Zhao D-L, Oosterhuis D (1997) Physiological response of growth chamber-grown cotton plants to the plant growth regulator PGR-IV under water-deficit stress. *Environ Exper Botany* 38: 7-14.
- Zhou Z-G, Meng Y-L, Shi Pei, Shen Y-Q, Jia Z-K (2000) Study of the relationship between boll weight in wheat-cotton double cropping and meteorological factors at boll-forming stage. *Acta Gossypii Sinica* 12: 122-126.






Figure 1. Daily number of flowers and bolls during the production stage (68 days) in the first season (I) for the Egyptian cotton cultivar Giza 75 (*Gossypium barbadense* L.) grown in uniform field trial at the experimental farm of the Agricultural Research Centre, Giza (30°N, 31°:28'E), Egypt. The soil texture was a clay loam, with an alluvial substratum, (pH = 8.07). Total water consumptive use during the growing season supplied by surface irrigation was about 6000 m³ha⁻¹. No rainfall occurred during the growing season. The sampling size was 261 plants (Sawan et al. 2005).




Figure 2. Daily number of flowers and bolls during the production stage (62 days) in the second season (II) for the Egyptian cotton cultivar Giza 75 (*Gossypium barbadense* L.) grown in uniform field trial at the experimental farm of the Agricultural Research Centre, Giza (30°N, 31°:28'E), Egypt. The soil texture was a clay loam, with an alluvial substratum, (pH = 8.07). Total water consumptive use during the growing season supplied by surface irrigation was about 6000 m³ha⁻¹. No rainfall occurred during the growing season. The sampling size was 358 plants (Sawan et al. 2005).




Figure 3. Daily boll retention ratio during the production stage (68 days) in the first season (I) for the Egyptian cotton cultivar Giza 75 (*Gossypium barbadense* L.) grown in uniform field trial at the experimental farm of the Agricultural Research Centre, Giza (30°N, 31°:28'E at an altitude 19 m), Egypt. The soil texture was a clay loam, with an alluvial substratum, (pH = 8.07). Total water consumptive use during the growing season supplied by surface irrigation was about 6000 m³ha⁻¹. No rainfall occurred during the growing season. The sampling size was 261 plants (Sawan et al. 2002).




Figure 4. Daily boll retention ratio during the production stage (62 days) in the second (II) for the Egyptian cotton cultivar Giza 75 (*Gossypium barbadense* L.) grown in uniform field trial at the experimental farm of the Agricultural Research Centre, Giza (30°N, 31°:28'E at an altitude 19 m), Egypt. The soil texture was a clay loam, with an alluvial substratum, (pH = 8.07). Total water consumptive use during the growing season supplied by surface irrigation was about 6000 m³ha⁻¹. No rainfall occurred during the growing season. The sampling size was 358 plants (Sawan et al. 2002).

Table 1. Range and mean values of the independent variables for the two seasons and over all data.

Climatic factor's	First season*		Second season**		Over all data (Two seasons)	
	Range	Mean	Range	Mean	Range	Mean
Max Temp (°C), (X ₁)	31.0-44.0	34.3	30.6-38.8	34.1	30.6-44.0	34.2
Min Temp (°C), (X ₂)	18.6-24.5	21.9	18.4-23.9	21.8	18.4-24.5	21.8
Max-Min Temp (°C), (X ₃) [®]	9.4-20.9	12.4	8.5-17.6	12.2	8.5-20.9	12.3
Evap (mm d ⁻¹), (X ₄)	7.6-15.2	10.0	4.1-9.8	6.0	4.1-15.2	8.0
0600 h Temp (°C), (X ₅)	14.0-21.5	17.8	13.3-22.4	18.0	13.3-22.4	17.9
1800 h Temp (°C), (X ₆)	19.6-27.0	24.0	20.6-27.4	24.2	19.6-27.4	24.1
Sunshine (h d ⁻¹), (X ₇)	10.3-12.9	11.7	9.7-13.0	11.9	9.7-13.0	11.8
Max RH (%), (X ₈)	62-96	85.4	51-84	73.2	51-96	79.6
Min RH (%), (X ₉)	11-45	30.8	23-52	39.8	11-52	35.1
Wind speed (m s ⁻¹), (X ₁₀)	ND	ND	2.2-7.8	4.6	ND	ND

(Sawan et al. 2006).

*Diurnal temperature range. ND not determined.

® Flower and boll stage (68 days, from 23 June through 29 August). ** Flower and boll stage (62 days, from 29 June through 29 August).

Table 2. Simple correlation values for the relationships between the independent variables and the studied dependent variable.

Independent variables (Climatic factors)	Dependent variable					
	First season		Second season		Combined data	
	Flower	Boll	Flower	Boll	Flower	Boll
Max Temp [°C] (X ₁)	-0.07	-0.03	-0.42**	-0.42**	-0.27**	-0.26**
Min Temp [°C] (X ₂)	-0.06	-0.07	0.00	0.02	-0.03	-0.02
Max-Min Temp [°C] (X ₃)	-0.03	-0.01	-0.36**	-0.37**	-0.25**	-0.24**
Evapor [mm d ⁻¹] (X ₄)	-0.56**	-0.53**	-0.61**	-0.59**	-0.40**	-0.48**
0600 h Temp [°C] (X ₅)	-0.01	-0.06	-0.14	-0.13	-0.09	-0.09
1800 h Temp [°C] (X ₆)	-0.02	-0.16	-0.37**	-0.36**	-0.27**	-0.25**
Sunshine [h d ⁻¹] (X ₇)	-0.25*	-0.14	-0.37**	-0.36**	-0.31**	-0.25**
Max RH [%] (X ₈)	0.40**	0.37**	0.01	0.01	0.04	-0.06
Min RH [%] (X ₉)	0.14	0.10	0.45**	0.46**	0.33**	0.39**
Wind speed [m s ⁻¹] (X ₁₀)	ND	ND	-0.06	-0.04	ND	ND

(Sawan et al. 2002). ND not determined * P < 0.05; ** P < 0.01.

Table 3. Selected factors and their relative contribution to variations of flower and boll production

Selected climatic factors	Flower production			Boll production		
	* R.C. (%)			R.C. (%)		
	First sea- son	Second season	Combined data	First sea- son	Second season	Combined data
Max Temp [°C] (X_1)	—	5.92	—	—	5.03	—
Evapor [mm d ⁻¹] (X_4)	19.08	23.45	16.06	23.04	22.39	22.89
1800 h Temp [°C] (X_6)	—	—	5.83	—	—	2.52
Sunshine [h d ⁻¹] (X_7)	9.43	7.77	8.31	11.65	7.88	5.47
Max RH [%] (X_8)	8.46	—	—	—	—	—
Min RH [%] (X_9)	—	4.37	7.38	—	4.26	4.64
** R ² % for selected factors	36.97	41.47	37.58	34.69	39.56	35.52
R ² % for factors studied	40.22	45.03	40.73	38.43	42.87	37.90
R ² % for factors deleted	3.25	3.56	3.15	3.74	3.31	2.38

(Sawan et al. 2002). * R.C. % = Relative contribution of each of the selected independent variables to variations of the dependent variable. ** R² % = Coefficient of determination in percentage form.

Table 4. Significant simple correlation coefficient values between the production variables and the studied climatic factors for the daily and different intervals of days combined over both seasons.

Daily and intervals of days	Production variables	Climatic factors ^z								
		Air temp (°C)			Evap (mm d ⁻¹)	Surface soil temp (°C)		Sunshine duration (h d ⁻¹)	Relative humidity (%)	
		Max (X_1)	Min (X_2)	Max-Min (X_3)	(X_4)	0600 h (X_5)	1800 h (X_6)	(X_7)	Max (X_8)	Min (X_9)
Daily (n = 120)	Flower	-0.15 ⁺⁺	NS	-0.26 ^{**}	-0.33 ^{**}	NS	-0.20 [*]	-0.23 [*]	NS	0.30 ^{**}
	Boll	NS	NS	-0.25 ^{**}	-0.43 ^{**}	NS	-0.19 ⁺⁺	-0.18 ⁺⁺	NS	0.36 ^{**}
	Boll ret. rat.	NS	NS	NS	-0.56 ^{**}	NS	NS	NS	NS	0.34 ^{**}
2 Days (n [#] = 60)	Flower	-0.31 ⁺⁺	NS	-0.32 [*]	-0.36 ^{**}	NS	-0.24 ⁺	-0.36 ^{**}	NS	0.37 ^{**}
	Boll	-0.29 ⁺⁺	NS	-0.30 ⁺⁺	-0.46 ^{**}	NS	-0.21 ⁺	-0.31 [*]	NS	0.44 ^{**}
	Boll ret. rat.	NS	NS	NS	-0.61 ^{**}	NS	NS	NS	NS	0.40 ^{**}
3 Days (n [#] = 40)	Flower	-0.34 [*]	NS	-0.34 [*]	-0.33 [*]	NS	-0.28 ⁺⁺	-0.39 [*]	NS	0.34 [*]
	Boll	-0.32 [*]	NS	-0.32 [*]	-0.48 ^{**}	NS	-0.24 ⁺	-0.36 [*]	NS	0.45 ^{**}
	Boll ret. rat.	NS	NS	NS	-0.63 ^{**}	NS	NS	NS	NS	0.40 [*]
4 Days (n [#] = 30)	Flower	-0.31 ⁺⁺	NS	-0.35 ⁺⁺	-0.33 ⁺⁺	NS	-0.28 ⁺	-0.39 [*]	NS	0.34 ⁺⁺
	Boll	-0.31 ⁺⁺	NS	-0.33 ⁺⁺	-0.48 ^{**}	NS	-0.23 ⁺	-0.38 [*]	NS	0.45 [*]
	Boll ret. rat.	NS	NS	NS	-0.64 ^{**}	NS	NS	NS	NS	0.42 [*]
5 Days (n [#] = 24)	Flower	-0.35 ⁺⁺	NS	-0.37 ⁺⁺	-0.39 ⁺⁺	NS	-0.39 ⁺⁺	-0.52 ^{**}	NS	0.41 [*]
	Boll	-0.33 ⁺	NS	-0.35 ⁺⁺	-0.49 [*]	NS	-0.35 ⁺⁺	-0.44 [*]	NS	0.47 ^{**}
	Boll ret. rat.	NS	NS	NS	-0.66 ^{**}	NS	NS	NS	NS	0.43 [*]
6 Days (n [#] = 20)	Flower	-0.37 ⁺⁺	NS	-0.41 ⁺⁺	-0.38 ⁺⁺	NS	NS	-0.54 ^{**}	NS	0.42 [*]
	Boll	-0.37 ⁺⁺	NS	-0.40 ⁺⁺	-0.49 [*]	NS	NS	-0.46 [*]	NS	0.49 [*]
	Boll ret. rat.	NS	NS	NS	-0.69 ^{**}	NS	NS	NS	NS	0.45 [*]
10 Days (n [#] = 12)	Flower	NS	NS	-0.45 ⁺⁺	-0.40 ⁺	NS	-0.55 [*]	-0.65 [*]	NS	0.43 ⁺⁺
	Boll	NS	NS	-0.43 ⁺⁺	-0.51 ⁺⁺	NS	-0.53 ⁺⁺	-0.57 [*]	NS	0.51 ⁺⁺
	Boll ret. rat.	NS	NS	NS	-0.74 ^{**}	NS	NS	NS	NS	0.55 [*]

(Sawan et al. 2006). ^z Wind speed did not show significant effect upon the studied production variables, so is not reported.

^{**} Significant at 1 % probability level, ^{*} Significant at 5 % probability level, ⁺⁺ Significant at 10 % probability level, ⁺ Significant at 15 % probability level. NS Means simple correlation coefficient is not significant at the 15% probability level. [#]n = Number of data pairs used in calculation.

Table 5. The equations obtained for each of the studied cotton production variables for the five-day intervals and daily intervals combined over both seasons.

Equation ^z	R ²	Significance
Five-day intervals		
$Y_1 = 23.78 - 0.5362X_4 - 0.1429X_6 - 0.1654X_7 + 0.0613X_9$	0.6237	**
$Y_2 = 15.89 - 0.4762X_4 - 0.1583X_6 - 0.1141X_7 + 0.0634X_9$	0.5945	**
$Y_3 = 72.65 - 0.0833X_4 - 0.1647X_6 + 0.2278X_9$	0.6126	**
Daily intervals		
$Y_1 = 19.78 - 0.181X_3 - 0.069X_4 - 0.164X_6 - 0.182X_7 + 0.010X_9$	0.4117	**
$Y_2 = 14.96 - 0.173X_3 - 0.075X_4 - 0.176X_6 - 0.129X_7 + 0.098X_9$	0.4461	**
$Y_3 = 52.36 - 3.601X_4 - 0.2352X_7 + 4.511X_9$	0.3587	**

(Sawan et al. 2006). ^zWhere Y_1 = number of flowers per plant, Y_2 = number of bolls per plant, Y_3 = boll retention ratio, X_3 = maximum – minimum temperature °C, X_4 = evaporation mm day⁻¹, X_6 = surface soil temperature °C at 1800 h., X_7 = sunshine duration h day⁻¹, and X_9 = minimum relative humidity %.

Table 6. Mean, standard deviation, maximum and minimum values of the climatic factors during the flower and boll stage (initial time) and the 15 days prior to flowering or subsequent to boll setting for I and II season at Giza, Egypt.

Climatic factors	First season*				Second season**			
	Mean	S.D.	Max.	Min.	Mean	S.D.	Max.	Min.
Max temp [°C] (X_1)	34.1	1.2	44.0	31.0	33.8	1.2	38.8	30.6
Min temp [°C] (X_2)	21.5	1.0	24.5	18.6	21.4	0.9	24.3	18.4
Max-Min temp [°C] (X_3) [□]	12.6	1.1	20.9	9.4	12.4	1.3	17.6	8.5
Evapor [mm d ⁻¹] (X_4)	10.6	1.6	16.4	7.6	6.0	0.7	9.8	4.1
0600 h temp [°C] (X_5)	17.5	1.1	21.5	13.9	17.6	1.2	22.4	13.3
1800 h temp [°C] (X_6)	24.2	1.9	32.3	19.6	23.7	1.1	27.4	20.6
Sunshine [h d ⁻¹] (X_7)	11.7	0.8	12.9	9.9	11.7	0.4	13.0	10.3
Max hum [%] (X_8)	85.6	3.3	96.0	62.0	72.9	3.8	84.0	51.0
Min hum [%] (X_9)	30.2	5.2	45.0	11.0	39.1	5.0	52.0	23.0
Wind speed [m s ⁻¹] (X_{10})	ND	ND	ND	ND	4.6	0.9	7.8	2.2

*Flower and boll stage (68 days, from 23 June through 29 August).

**Flower and boll stage (62 days, from 29 June through 29 August). (Sawan et al. 2005).

Table 7. Simple correlation coefficients (r) between climatic factors and number of flower and harvested bolls in initial time (0) and each of the 15-day periods before flowering in the first season (I).

Climate period		Air temp. (°C)			Evap. (mm d ⁻¹)	Surface soil temp. (°C)		Sunshine duration (h d ⁻¹)	Humidity (%)	
		Max. (X ₁)	Min. (X ₂)	Max-Min [♦] (X ₃)	(X ₄)	0600 h (X ₅)	1800 h (X ₆)	(X ₇)	Max. (X ₈)	Min. (X ₉)
0 [#]	Flower	-0.07	-0.06	-0.03	-0.56**	-0.01	-0.20	-0.25*	0.40**	0.14
	Boll	-0.03	-0.07	-0.01	-0.53**	-0.06	-0.16	-0.14	0.37**	0.10
1	Flower	-0.15	-0.08	-0.11	-0.64**	-0.01	-0.17	-0.30*	0.39**	0.20
	Boll	-0.07	-0.08	-0.02	-0.58**	-0.06	-0.10	-0.23*	0.36**	0.13
2	Flower	-0.26*	-0.10	-0.22	-0.69**	-0.07	-0.30*	-0.35**	0.42**	0.30*
	Boll	-0.18	-0.08	-0.14	-0.64**	-0.05	-0.21	-0.25*	0.40**	0.20
3	Flower	-0.28*	-0.02	-0.31**	-0.72**	0.15	-0.29*	-0.37**	0.46**	0.35**
	Boll	-0.19	-0.02	-0.21	-0.65**	0.11	-0.20	-0.30*	0.37**	0.25*
4	Flower	-0.26*	-0.03	-0.26*	-0.67**	0.08	-0.24*	-0.41**	0.46**	0.35**
	Boll	-0.21	-0.04	-0.21	-0.63**	0.04	-0.18	-0.35**	0.39**	0.29*
5	Flower	-0.27*	-0.02	-0.27*	-0.68**	0.16	-0.29*	-0.45**	0.49**	0.38**
	Boll	-0.22	0.00	-0.24*	-0.63**	0.16	-0.21	-0.39**	0.44**	0.32**
6	Flower	-0.21	0.05	-0.25*	-0.73**	0.16	-0.28*	-0.46**	0.47**	0.42**
	Boll	-0.15	0.08	-0.21	-0.67**	0.19	-0.19	-0.46**	0.43**	0.35**
7	Flower	-0.17	-0.01	-0.17	-0.69**	0.10	-0.27*	-0.43**	0.46**	0.35**
	Boll	-0.11	-0.06	-0.15	-0.64**	0.14	-0.19	-0.46**	0.43**	0.32**
8	Flower	-0.24*	-0.03	-0.24*	-0.71**	0.09	-0.30*	-0.44**	0.45**	0.45**
	Boll	-0.14	0.04	-0.17	-0.63**	0.16	-0.17	-0.48**	0.44**	0.39**
9	Flower	-0.23	-0.10	-0.19	-0.68**	0.05	-0.33**	-0.32**	0.43**	0.44**
	Boll	-0.14	0.04	-0.17	-0.61**	0.15	-0.21	-0.40**	0.42**	0.41**
10	Flower	-0.26*	0.05	-0.30*	-0.67**	0.13	-0.29*	-0.29*	0.40**	0.48**
	Boll	-0.14	0.13	-0.22	-0.58**	0.22	-0.17	-0.36**	0.46**	0.41**
11	Flower	-0.20	0.10	-0.27*	-0.62**	0.21	-0.19	-0.29*	0.42**	0.44**
	Boll	-0.04	0.22	-0.16	-0.53**	0.27*	-0.04	-0.38**	0.45**	0.36**
12	Flower	-0.17	0.16	-0.26*	-0.62**	0.29*	-0.15	-0.40**	0.44**	0.45**
	Boll	0.00	0.25*	-0.13	-0.51**	0.35**	-0.04	-0.45**	0.40**	0.30*
13	Flower	-0.13	0.16	-0.22	-0.62**	0.23	-0.12	-0.42**	0.43**	0.45**
	Boll	0.00	0.22	-0.11	-0.51**	0.30*	-0.03	-0.49**	0.41**	0.33**
14	Flower	-0.08	0.18	-0.18	-0.56**	0.21	-0.15	-0.44**	0.41**	0.46**
	Boll	0.01	0.21	-0.10	-0.47**	0.26*	-0.09	-0.49**	0.42**	0.33**
15	Flower	-0.08	0.22	-0.21	-0.51**	0.24*	-0.22	-0.42**	0.39**	0.38**
	Boll	-0.03	0.19	-0.13	-0.45**	0.24*	-0.17	-0.44**	0.43**	0.30*

*: Significant at 5% level and **: significant at 1% level.

0 = Initial time.

♦ diurnal temperature range.

(Sawan et al. 2005).

Table 8. Simple correlation coefficients (r) between climatic factors^z and number of flower and harvested bolls in initial time (0) and each of the 15-day periods before flowering in the second season (II).

Climate period		Air temp. (°C)			Evap. (mm d ⁻¹)	Surface soil temp. (°C)		Sunshine duration (h d ⁻¹)	Humidity (%)	
		Max. (X ₁)	Min. (X ₂)	Max-Min [♦] (X ₃)	(X ₄)	0600 h (X ₅)	1800 h (X ₆)	(X ₇)	Max. (X ₈)	Min. (X ₉)
0 [#]	Flower	-0.42**	0.00	-0.36**	-0.61**	-0.14	-0.37**	-0.37**	0.01	0.45**
	Boll	-0.42**	0.02	-0.37**	-0.59**	-0.13	-0.36**	-0.36**	0.01	0.46**
1	Flower	-0.42**	0.10	-0.42**	-0.63**	-0.08	-0.29*	-0.41**	0.05	0.48**
	Boll	-0.41**	0.11	-0.42**	-0.62**	-0.07	-0.28*	-0.41**	0.05	0.47**
2	Flower	-0.40**	0.08	-0.43**	-0.65**	-0.09	-0.27*	-0.39**	0.02	0.49**
	Boll	-0.40**	0.08	-0.43**	-0.64**	-0.08	-0.26*	-0.40**	0.03	0.49**
3	Flower	-0.38**	0.13	-0.43**	-0.61**	-0.06	-0.17	-0.38**	0.00	0.45**
	Boll	-0.37**	0.15	-0.44**	-0.61**	-0.05	-0.15	-0.38**	0.01	0.46**
4	Flower	-0.36**	0.17	-0.41**	-0.61**	-0.04	-0.18	-0.38**	0.02	0.45**
	Boll	-0.35**	0.18	-0.41**	-0.60**	-0.03	-0.16	-0.36**	0.03	0.44**
5	Flower	-0.30*	0.13	-0.36**	-0.60**	-0.07	-0.23	-0.32**	-0.05	0.43**
	Boll	-0.28*	0.15	-0.35**	-0.58**	-0.05	-0.21	-0.31**	-0.05	0.41**
6	Flower	-0.24	0.21	-0.38**	-0.61**	-0.02	-0.12	-0.28*	0.02	0.40**
	Boll	-0.22	0.24	-0.38**	-0.59**	0.00	-0.07	-0.29*	0.02	0.40**
7	Flower	-0.19	0.23	-0.29*	-0.54**	-0.03	-0.05	-0.26*	-0.04	0.32**
	Boll	-0.18	0.23	-0.27*	-0.53**	-0.02	-0.03	-0.27*	-0.04	0.30*
8	Flower	-0.15	0.24	-0.25*	-0.52**	-0.03	-0.07	-0.24*	-0.05	0.28*
	Boll	-0.14	0.22	-0.22	-0.51**	-0.03	-0.06	-0.22*	-0.05	0.26*
9	Flower	-0.16	0.34**	-0.32**	-0.56**	0.08	-0.02	-0.25*	0.05	0.30*
	Boll	-0.14	0.34**	-0.31**	-0.56**	0.09	-0.01	-0.23*	0.07	0.29*
10	Flower	-0.16	0.31**	-0.30*	-0.56**	0.11	-0.06	-0.27*	0.11	0.33**
	Boll	-0.14	0.28*	-0.27*	-0.55**	0.09	-0.07	-0.25*	0.09	0.31**
11	Flower	-0.16	0.31**	-0.27*	-0.55**	0.10	-0.02	-0.31**	0.08	0.32**
	Boll	-0.15	0.29*	-0.26*	-0.53**	0.10	0.00	-0.29*	0.08	0.29*
12	Flower	-0.17	0.44**	-0.37**	-0.57**	0.26*	0.02	-0.36**	0.17	0.34**
	Boll	-0.17	0.42**	-0.36**	-0.55**	0.25*	0.01	-0.34**	0.16	0.32**
13	Flower	-0.14	0.40**	-0.33**	-0.56**	0.21	0.03	-0.28*	0.10	0.34**
	Boll	-0.15	0.38**	-0.34**	-0.56**	0.21	0.01	-0.27*	0.09	0.33**
14	Flower	-0.19	0.39**	-0.38**	-0.59**	0.25*	0.04	-0.34**	0.16	0.35**
	Boll	-0.20	0.39**	-0.40**	-0.59**	0.26*	0.03	-0.36**	0.17	0.36**
15	Flower	-0.24	0.49**	-0.45**	-0.62**	0.37**	0.16	-0.38**	0.27*	0.42**
	Boll	-0.24	0.51**	-0.48**	-0.63**	0.40**	0.15	-0.40**	0.26*	0.43**

*: Significant at 5% level and **: significant at 1% level.

[#] 0 = Initial time.[♦] diurnal temperature range.^z Wind speed did not show significant effect upon the studied production variables, so it is not reported. (Sawan et al. 2005).

Table 9. Simple correlation coefficient (r) values between climatic factors and number of harvested bolls and retention ratio in initial time (0) and each of the 15-day periods after flowering in the first season (I).

Climate period		Air temp. (°C)			Evap. (mm d ⁻¹)	Surface soil temp. (°C)		Sunshine duration (h d ⁻¹)	Humidity (%)	
		Max. (X ₁)	Min. (X ₂)	Max.-Min.* (X ₃)	(X ₄)	0600 h (X ₅)	1800 h (X ₆)	(X ₇)	Max. (X ₈)	Min. (X ₉)
0 [#]	Retention ratio•	-0.05	-0.03	-0.03	-0.10	-0.11	0.10	0.20	-0.04	-0.02
	No. of bolls	-0.03	-0.07	-0.01	-0.53**	-0.06	-0.16	-0.14	0.37**	0.10
1	Retention ratio	-0.07	-0.08	-0.01	-0.10	-0.16	0.04	0.15	0.04	0.05
	No. of bolls	0.02	-0.08	0.08	-0.49**	-0.09	-0.05	-0.20	0.35**	0.09
2	Retention ratio	-0.08	-0.14	0.02	-0.08	-0.19	0.03	0.17	0.02	-0.02
	No. of bolls	0.02	-0.04	0.07	-0.46**	-0.06	-0.01	-0.19	0.33**	0.09
3	Retention ratio	-0.09	-0.21	0.06	-0.08	-0.24*	0.02	0.19	0.01	-0.10
	No. of bolls	0.03	-0.03	0.06	-0.44**	-0.04	0.05	-0.18	0.32**	0.08
4	Retention ratio	-0.05	-0.20	0.09	-0.01	-0.24*	0.01	0.22	0.00	-0.15
	No. of bolls	0.01	-0.05	0.05	-0.40**	-0.03	0.04	-0.16	0.31*	0.08
5	Retention ratio	-0.03	-0.21	0.13	0.07	-0.25*	0.00	0.26*	-0.02	-0.22
	No. of bolls	0.00	-0.07	0.05	-0.37**	-0.02	0.03	-0.13	0.29*	0.07
6	Retention ratio	0.01	-0.19	0.15	0.12	-0.24*	0.02	0.27*	-0.03	-0.20
	No. of bolls	-0.01	-0.08	0.04	-0.38**	-0.02	0.04	-0.15	0.31*	0.13
7	Retention ratio	0.05	-0.17	0.17	0.18	-0.25*	0.05	0.29*	-0.02	-0.21
	No. of bolls	-0.03	-0.09	0.03	-0.39**	-0.04	0.06	-0.14	0.34**	0.18
8	Retention ratio	0.06	-0.08	0.13	0.21	-0.20	0.07	0.28*	-0.06	-0.19
	No. of bolls	-0.05	-0.07	-0.01	-0.35**	-0.02	0.02	-0.17	0.28*	0.17
9	Retention ratio	0.08	0.00	0.08	0.26*	-0.14	0.08	0.29*	-0.12	-0.20
	No. of bolls	-0.08	-0.06	-0.05	-0.33**	-0.01	0.00	-0.23	0.20	0.16
10	Retention ratio	0.06	-0.02	0.05	0.27*	-0.13	0.09	0.27*	-0.10	-0.08
	No. of bolls	-0.11	-0.10	-0.07	-0.34**	-0.03	-0.03	-0.19	0.18	0.21
11	Retention ratio	0.04	-0.04	0.08	0.28*	-0.12	0.08	0.26*	-0.09	-0.05
	No. of bolls	-0.18	-0.18	-0.06	-0.37**	-0.10	-0.04	-0.14	0.15	
0.28*										
12	Retention ratio	0.02	0.01	-0.08	0.32**	-0.05	0.05	0.25*	-0.08	-0.03
	No. of bolls	-0.17	-0.13	-0.08	-0.32**	-0.06	-0.07	-0.11	0.16	
0.24*										
13	Retention ratio	-0.04	0.04	-0.09	0.38**	0.00	0.01	0.27*	-0.09	-0.02
	No. of bolls	-0.15	-0.09	-0.09	-0.29*	-0.03	-0.10	-0.08	0.18	0.20
14	Retention ratio	-0.07	0.04	-0.13	0.34**	0.06	-0.02	0.18	-0.08	-0.01
	No. of bolls	-0.15	-0.10	-0.10	-0.28*	-0.01	-0.10	-0.15	0.17	0.17
15	Retention ratio	-0.13	0.03	-0.18	0.33**	0.09	-0.04	0.06	-0.07	0.00
	No. of bolls	-0.16	-0.10	-0.11	-0.28*	0.00	-0.11	-0.13	0.17	0.15

* and ** Significant at 5% and 1% levels of significance, respectively. [#] 0 = Initial time • Retention ratio: (the number of retained bolls obtained from the total number of each daily tagged flowers in all selected plants at harvest/each daily number of tagged flowers in all selected plants) x 100. ♦ diurnal temperature range. (Sawan et al. 2005).

Table 10. Simple correlation coefficient (r) values between climatic factors^z and number of harvested bolls and retention ratio in initial time (0) and each of the 15-day periods after flowering in the second season (II).

Climate period		Air temp. (°C)			Evap. (mm d ⁻¹)	Surface soil temp. (°C)		Sunshine duration (h d ⁻¹)	Humidity (%)	
		Max. (X ₁)	Min. (X ₂)	Max.-Min [♦] (X ₃)		0600 h (X ₅)	1800 h (X ₆)		Max. (X ₈)	Min. (X ₉)
0 [#]	Retention ratio•	-0.04	0.20	-0.31*	-0.14	0.12	-0.20	0.01	-0.04	0.17
	No. of bolls	-0.42**	0.02	-0.37**	-0.59**	-0.13	-0.36**	-0.36**	0.01	0.46**
1	Retention ratio	-0.10	-0.03	-0.22	-0.21	-0.15	-0.05	-0.04	-0.02	0.23
	No. of bolls	-0.25*	-0.01	-0.36**	-0.63**	-0.15	-0.30*	-0.25*	0.06	0.44**
2	Retention ratio	-0.15	-0.06	-0.10	-0.15	-0.08	-0.21	-0.01	-0.04	0.12
	No. of bolls	-0.18	-0.01	-0.34**	-0.65**	-0.11	-0.25*	-0.32*	0.13	0.43**
3	Retention ratio	-0.03	-0.01	-0.02	-0.21	-0.01	-0.17	-0.08	0.09	0.12
	No. of bolls	-0.15	-0.06	-0.30*	-0.62**	-0.05	-0.28*	-0.31*	0.14	0.33**
4	Retention ratio	0.08	-0.02	0.07	-0.09	-0.03	-0.09	-0.10	0.05	-0.04
	No. of bolls	-0.15	-0.05	-0.28*	-0.63**	-0.06	-0.25*	-0.33**	0.15	0.32*
5	Retention ratio	0.23	-0.03	0.12	-0.06	-0.06	-0.01	-0.11	0.01	-0.16
	No. of bolls	-0.14	-0.05	-0.25*	-0.62**	-0.06	-0.24*	-0.35**	0.15	0.31*
6	Retention ratio	0.09	-0.08	0.12	-0.09	-0.07	-0.01	-0.09	0.00	-0.05
	No. of bolls	-0.15	-0.04	-0.22	-0.61**	-0.08	-0.25*	-0.34**	0.13	0.22
7	Retention ratio	-0.03	-0.12	0.12	-0.10	-0.11	-0.01	-0.04	-0.03	0.02
	No. of bolls	-0.15	-0.02	-0.19	-0.60**	-0.10	-0.29*	-0.32*	0.10	0.18
8	Retention ratio	-0.02	0.05	0.03	-0.10	-0.04	-0.03	-0.02	-0.01	0.01
	No. of bolls	-0.20	-0.03	-0.23	-0.61**	-0.10	-0.28*	-0.32*	0.19	0.22
9	Retention ratio	-0.02	0.13	-0.05	-0.10	0.08	-0.05	-0.01	0.03	0.00
	No. of bolls	-0.24	-0.04	-0.29*	-0.62**	-0.11	-0.30*	-0.33**	0.13	0.27*
10	Retention ratio	-0.04	0.12	-0.08	-0.09	0.05	0.11	-0.02	0.04	0.02
	No. of bolls	-0.27*	-0.07	-0.30*	-0.60**	-0.16	-0.34**	-0.34**	0.11	0.26*
11	Retention ratio	-0.07	0.10	-0.10	-0.08	0.03	0.20	-0.03	0.05	0.04
	No. of bolls	-0.30*	-0.12	-0.30*	-0.61**	-0.18	-0.39**	-0.36**	0.10	0.27*
12	Retention ratio	-0.11	0.09	-0.14	-0.11	0.04	0.13	-0.08	0.11	0.09
	No. of bolls	-0.32*	-0.19	-0.26*	-0.60**	-0.22	-0.42**	-0.37**	0.09	0.27*
13	Retention ratio	-0.14	0.09	-0.17	-0.18	0.06	-0.06	-0.14	0.16	0.12
	No. of bolls	-0.33**	-0.26*	-0.23	-0.59**	-0.28*	-0.48**	-0.39**	0.08	0.27*
14	Retention ratio	-0.11	-0.04	-0.10	-0.13	-0.15	-0.05	-0.09	0.01	0.12
	No. of bolls	-0.34**	-0.32*	-0.21	-0.61**	-0.32*	-0.48**	-0.38**	0.06	0.27*
15	Retention ratio	-0.08	-0.11	0.02	-0.08	-0.22	-0.05	-0.02	-0.03	0.12
	No. of bolls	-0.35**	-0.37**	-0.18	-0.61**	-0.38**	-0.48**	-0.37**	0.03	0.27*

* and ** Significant at 5% and 1% levels of significance, respectively.

[#] 0 = Initial time

• Retention ratio: (the number of retained bolls obtained from the total number of each daily tagged flowers in all selected plants at harvest/each daily number of tagged flowers in all selected plants) x 100.

♦ diurnal temperature range.

^z Wind speed did not show significant effect upon the studied production variables, so it is not reported.

(Sawan et al. 2005).

Table 11. The models obtained for the number of flowers and bolls per plant as functions of the climatic data derived from the 5, 10, and 15 day periods prior to flower opening in the two seasons (I, II).

Season	Model ^z	R ²	Significance
First			
<i>Flower</i>			
	$Y_1 = 55.75 + 0.86X_3 - 2.09X_4 - 2.23X_7$	0.51	**
	$Y_2 = 26.76 - 5.45X_4 + 1.76X_9$	0.42	**
	$Y_3 = 43.37 - 1.02X_4 - 2.61X_7 + 0.20X_8$	0.52	**
<i>Boll</i>			
	$Y_1 = 43.69 + 0.34X_3 - 1.71X_4 - 1.44X_7$	0.43	**
	$Y_2 = 40.11 - 1.82X_4 - 1.36X_7 + 0.10X_8$	0.48	**
	$Y_3 = 31.00 - 0.60X_4 - 2.62X_7 + 0.23X_8$	0.47	**
Second			
<i>Flower</i>			
	$Y_1 = 18.58 + 0.39X_3 - 0.22X_4 - 1.19X_7 + 0.17X_9$	0.54	**
	$Y_2 = 16.21 + 0.63X_3 - 0.20X_4 - 1.24X_7 + 0.16X_9$	0.61	**
	$Y_3 = 14.72 + 0.51X_3 - 0.20X_4 - 0.85X_7 + 0.17X_9$	0.58	**
<i>Boll</i>			
	$Y_1 = 25.83 + 0.50X_3 - 0.26X_4 - 1.95X_7 + 0.15X_9$	0.61	**
	$Y_2 = 19.65 + 0.62X_3 - 0.25X_4 - 1.44X_7 + 0.12X_9$	0.60	**
	$Y_3 = 15.83 + 0.60X_3 - 0.22X_4 - 1.26X_7 + 0.14X_9$	0.59	**

^z Where Y_1 , Y_2 , Y_3 = number of flowers or bolls per plant at the 5, 10 and 15 day periods before flowering, respectively, X_2 = minimum temperature (°C), X_3 = diurnal temperature range (°C), X_4 = evaporation (mm day⁻¹), X_7 = sunshine duration (h day⁻¹), X_8 = maximum humidity (%) and X_9 = minimum humidity (%).

(Sawan et al. 2005).

Table 12. The models obtained for the number of bolls per plant as functions of the climatic data derived from the 5, 10, and 15 day periods after flower opening in the two seasons (I, II).

Season	Model ^z	R ²	Significance
First	$Y_1 = 16.38 - 0.41X_4$	0.14	**
	$Y_2 = 16.43 - 0.41X_4$	0.14	**
	$Y_3 = 27.83 - 0.60X_4 - 0.88X_9$	0.15	**
Second	$Y_1 = 23.96 - 0.47X_4 - 0.77X_8$	0.44	**
	$Y_2 = 18.72 - 0.58X_4$	0.34	**
	$Y_3 = 56.09 - 2.51X_4 - 0.49X_6 - 1.67X_7$	0.56	**

^zWhere Y_1 , Y_2 , Y_3 = number of bolls per plant at the 5, 10, and 15 day periods after flowering, respectively, X_4 = evaporation (mm day⁻¹), X_6 = soil surface temperature (°C) at 1800, X_7 = sunshine duration (h day⁻¹), X_8 = maximum humidity (%) and X_9 = minimum humidity (%).

(Sawan et al. 2005).

Table 13. Simple correlation coefficient (r) values between the independent variables and the dependent variables in the first season (I).

Independent variables (Irrigation and climatic factors)	Dependent variables (First season)	
	Flowers	Bolls
(X1) Irrigation on day 1	-0.1282	-0.0925
(X2) Irrigation on day 0 or -1 (1 st and 2 nd day after irrigation)	-0.1644	-0.1403
(X3) 1 is for the day prior to irrigation	-0.0891	-0.0897
(X4) Number of days that temperature equaled or exceeded 37.5 °C	0.1258	0.1525
(X5) Range of temperature [°C] on day 1	-0.0270	-0.0205
(X6) Broadest range of temperature [°C] over days 1 -12	0.0550	0.1788 ^d
(X7) MinRH [%] during day 1	0.1492	0.1167
(X8) MaxRH [%] during day 1	0.2087 ^c	0.1531
(X9) MinRH [%] during day 2	0.1079	0.1033
(X10) MaxRH [%] during day 2	0.1127	0.0455
(X11) Largest maxRH [%] on days 3-6	0.3905 ^a	0.2819 ^b
(X12) Lowest minRH [%] on days 3-6	0.0646	0.0444
(X13) Largest maxRH [%] on days 7-12	0.4499 ^a	0.3554 ^b
(X14) Lowest minRH [%] on days 7-12	0.3522 ^a	0.1937 ^d
(X15) Lowest minRH [%] on days 50-52	-0.3440 ^a	-0.4222 ^a
(X16) Daily light period (hour)	-0.2430 ^b	-0.1426

(Sawan et al. 2010). ^aSignificant at 1 % probability level ^bSignificant at 5 % probability level ^c Significant at 10 % probability level ^d Significant at 15 % probability level

Table 14. Simple correlation coefficient (r) values between the independent variables and the dependent variables in the second season (II).

Independent variables (Irrigation and climatic factors)	Dependent variables (Second season)	
	Flowers	Bolls
(X1) Irrigation on day 1	-0.0536	-0.0467
(X2) Irrigation on day 0 or -1	-0.1116	-0.1208
(X3) 1 is for the day prior to the day of irrigation	-0.0929	-0.0927
(X4) Number of days that temperature equaled or exceeded 37.5 °C	-0.4192 ^a	-0.3981 ^a
(X5) Range of temperature [°C] on day 1	-0.3779 ^a	-0.3858 ^a
(X6) Broadest range of temperature [°C] over days 1-12	-0.3849 ^a	-0.3841 ^a
(X7) MinRH [%] during day 1	0.4522 ^a	0.4665 ^a
(X8) MaxRH [%] during day 1	0.0083	0.0054
(X9) MinRH [%] during day 2	0.4315 ^a	0.4374 ^a
(X10) MaxRH [%] during day 2	0.0605	0.0532
(X11) Largest maxRH [%] on days 3-6	0.2486 ^c	0.2520 ^b
(X12) Lowest minRH [%] on days 3-6	0.5783 ^a	0.5677 ^a
(X13) Largest maxRH [%] on days 7-12	0.0617	0.0735
(X14) Lowest minRH [%] on days 7-12	0.4887 ^a	0.4691 ^a
(X15) Lowest minRH [%] on days 50-52	-0.6246 ^a	-0.6113 ^a
(X16) Daily light period (hour)	-0.3677 ^a	-0.3609 ^a

(Sawan et al. 2010). ^a Significant at 1 % probability level ^b Significant at 5 % probability level ^c Significant at 10 % probability level

Table 15. Simple correlation coefficient (r) values between the independent variables and dependent variables in the combined two seasons (I and II).

Independent variables (Irrigation and climatic factors)	Dependent variables (Combined two seasons)	
	Flowers	Bolls
(X1) Irrigation on day 1	-0.0718	-0.0483
(X2) Irrigation on day 0 or -1	-0.1214	-0.1108
(X3) 1 is for the day prior to the day of irrigation	-0.0845	-0.0769
(X4) Number of days that temperature equaled or exceeded 37.5 °C	-0.2234 ^b	-0.1720 ^c
(X5) Range of temperature [°C] on day 1	-0.2551 ^a	-0.2479 ^a
(X6) Broadest range of temperature [°C] over days 1-12	-0.2372 ^a	-0.1958 ^b
(X7) MinRH [%] during day 1	0.3369 ^a	0.3934 ^a
(X8) MaxRH [%] during day 1	0.0032	-0.0911
(X9) MinRH [%] during day 2	0.3147 ^a	0.3815 ^a
(X10) MaxRH [%] during day 2	-0.0094	-0.1113
(X11) Largest maxRH [%] on days 3-6	0.0606	-0.0663
(X12) Lowest minRH [%] on days 3-6	0.3849 ^a	0.4347 ^a
(X13) Largest maxRH [%] on days 7-12	-0.0169	-0.1442 ^d
(X14) Lowest minRH [%] on days 7-12	0.3891 ^a	0.4219 ^a
(X15) Lowest minRH [%] on days 50-52	-0.3035 ^a	-0.2359 ^a
(X16) Daily light period (hour)	-0.3039 ^a	-0.2535 ^a

(Sawan et al. 2010). ^a Significant at 1 % probability level ^b Significant at 5 % probability level ^c Significant at 10 % probability level ^d Significant at 15 % probability level

Table 16. Model obtained for cotton production variables as functions of climatic data and soil moisture status in individual and combined seasons.

Season	Model	R ²
Season I (n = 68)	$Y_1 = -557.54 + 6.35X_6 + 0.65X_7 + 1.92X_{11} + 4.17X_{13} + 2.88X_{14} - 1.90X_{15} - 5.63X_{16}$	0.63
	$Y_2 = -453.93 + 6.53X_6 + 0.61X_7 + 1.80X_{11} + 2.47X_{13} + 1.87X_{14} - 1.85X_{15}$	0.53
Season II (n = 62)	$Y_1 = -129.45 + 25.36X_1 + 37.02X_4 + 1.48X_7 + 1.69X_9 + 4.46X_{12} + 2.55X_{14} - 4.73X_{15}$	0.72
	$Y_2 = -130.23 + 24.27X_1 + 35.66X_4 + 1.42X_7 + 1.61X_9 + 4.00X_{12} + 2.18X_{14} - 4.09X_{15}$	0.71
Combined data: I & II (n = 130)	$Y_1 = -557.36 + 6.82X_6 + 1.44X_7 + 0.75X_9 + 2.04X_{11} + 2.55X_{12} + 2.01X_{13} + 3.27X_{14} - 2.15X_{15}$	0.57
	$Y_2 = -322.17 + 6.41X_6 + 1.20X_7 + 0.69X_9 + 1.81X_{11} + 2.12X_{12} + 2.35X_{14} - 2.16X_{15}$	0.53

(Sawan et al. 2010).

(Y1) Number of cotton flowers; (Y2) Number of cotton bolls.

(X1) Irrigation on day 1; (X4) Number of that temperature equaled or exceeded 37.5 °C; (X6) Broadest range of temperature [°C] over days 1-12; (X7) MinRH [%] during day 1; (X9) MinRH [%] during day 2; (X11) Largest maxRH [%] on days 3-6; (X12) Lowest minRH [%] on days 3-6; (X13) Largest maxRH [%] on days 7-12; (X14) Lowest minRH [%] on days 7-12; (X15) Lowest minRH [%] on days 50-52; (X16) Daily light period (hour).

All entries significant at 1% level.