



American Journal of Clinical Research and Reviews (ISSN:2576-0505)



The Mystery of Patient Behavior

AK Mohiuddin

Department of Pharmacy, World University of Bangladesh, Bangladesh

ABSTRACT

Patient behavior is often described as disruptive behavior as they have an altered mental stage of fear of being sick, anxious about out of the pocket cost, alteration of lifestyle if suffered from a chronic illness. And the outcomes often faced by providers are inappropriate language, make unreasonable demands, and may even resort to physical abuse. The article comprises a few of this unusual behavior and a simple comparison between patients from developed world with those of the under-developed east. Purpose of the study: Discussion and projection of behavior pattern, health seeking behavior and monitoring status in both developed and under-privileged countries. The pharmacists have a vital role to play which is discussed along with comparison. Findings: Developed or under developed country, patients have a separate behavior pattern which develops and worsen with disease progression mostly. So many factors are behind but one thing clearly understood that the handling of such situation is a provider's function, a challenge they have to face along with treatment intervention. Research Limitations: Very few articles found in matters regarding along with a very less interest paid by general people to talk about healthcare matters. It was very difficult to bring out facts of irrational patient behavior, giving it a substantial figure to discuss in this article. However, the major limitation is the article could be a comparison of behaviors of developed and under-privileged countries which requires an enormous exposure and financial support. However, the sole focus was to detail mysterious patient behavior and a greater part is covered. Practical Implication: The soul of this article was to detail about patient behavior, both in Bangladesh and developed countries. Along with students, researchers and professionals of different background and disciplines, eg. Pharmacists, marketers, doctors, nurses, hospital authorities, public representatives, policy makers and regulatory authorities have to acquire much from this article. Social Implication: Patient behavior is the Sole of healing system and a scope for pharmacists and other healthcare providers to work with in both developed or under developed countries. The article should contribute an integrated guideline for patient compliance, demand rational behavior and last but not the least a silvery lining to better pharmacists' dealings with them in near future.

Keywords:

Health Beliefs and Perception; Patient Counseling; Patient Comprehension and Recall; Patient Motivation; Patient Feedback; Patient Compliance.

*Correspondence to Author:

AK Mohiuddin

Department of Pharmacy, World University of Bangladesh, Bangladesh

How to cite this article:

AK Mohiuddin. The Mystery of Patient Behavior. American Journal of Clinical Research and Reviews, 2018; 3:8.



eSciPub LLC, Houston, TX USA.

Website: <http://escipub.com/>

1. Introduction

The unusual patient behavior toward providers and towards themselves including medication taking is inevitable. An interesting fact is mostly patients avoid going to visit a doctor or pharmacists and when appeared for treatment their behaviors change dramatically which is still unexplained. A National Health Service survey reveals 30% patients never seek any care due to emotional distress of being ill and again nearly 60% fail to comply treatment guidelines as directed. Social phobia persists in women who need to go for an intervention for any gynecological disorder. Conversely children have phobia with the doctors and hospital settings due to some unknown reason. When came late at an advanced disease state, the patient is already in a highly disturbed mental condition and a few situations raise, providers need to face another challenge to tolerate and mitigate them.

2. Materials and Methods

2.1. Research Design and Search Strategy

The research is conducted through secondary data search from several sources from books, technical newsletters, newspapers, journals, and many other sources. The present study was started from the beginning of 2018. PubMed, ALTAVISTA, Embase, Scopus, Web of Science, and the Cochrane Central Register of was thoroughly searched. The keywords were used to search for different publishers' journals like Elsevier, Springer, Willey Online Library, Wolters Kluwer were extensively followed.

2.2. Inclusion and Exclusion Criteria

Studies regarding psychological and social behaviors were given priorities. Several factors that influence medication taking behavior was collectively analyzed and added to the article. Issues regarding economic and cultural barriers were found to be different from subcontinents, countries and even states. A few interesting features of patient motivation and satisfaction through pharmacist's intervention were added afterwards to maintain a logical sequence. Drug

factors, environmental factors and provider-patient interaction followed by pharmacist's role in handling patients and to change or correction of behavior were added to reveal their effect on patient compliance which is the ultimate goal of therapeutic guidelines. Many studies found regarding patient compliance of different health conditions in different countries found to be not within the scope of this article.

3. Results and Discussion

3.1. Types of Medication Taking Behavior

- a) *Compliance*: Understanding how medication should be used and its benefit, with sufficient positive motivation, intentions and act accordingly.
- b) *Adherence*: The extent to which a person takes medication as prescribed. Adherence is broadly viewed as related to instructions concerning medicine intake, use of medical device, diet, exercise, life style changes, rest and return for scheduled appointments.
- c) *Concordance*: A process by which a patient and clinician make decisions together about treatment.
- d) *Persistence*: A person's ability to continue medical advice for the intended course, which may range from few days to life-long [1].

3.2. Types of patient behavior

a). *Health behavior* that is preventive in nature generally is referred to as preventive health behavior. Expanding on the original definition, preventive health behavior is defined as actions taken to prevent illness and maintain physical, emotional, intellectual, spiritual, and social well-being [2]. Examples of preventive health behaviors include

- Participation in health screening programs
- Following healthy diet recommendations
- Participation in relaxation and cardiovascular exercises
- Creating and maintaining close personal relationships

b) *Illness behavior* is any activity undertaken by individuals who perceive themselves to be ill that

defines the state of their health and aids in discovering a suitable remedy [3].

- Illness behavior is the way persons respond to bodily indications that they experience as abnormal; thus, it involves the manner in which persons monitor their bodies, define and interpret their symptoms, and seek health care.
- Individuals attempt to ascribe cause and meaning to their illness symptoms and may self-diagnose and treat.
- Alternatively, individuals may visit a doctor or another prescriber and a pharmacist in order to obtain a prescription drug.

c) *Treatment behavior* is actions taken to restore health or halt disease progression traditionally have been referred to as sick-role behaviors and now are referred to as treatment behaviors [4]. Model of Health Services Utilization suggested that three main factors affect an individual's use of health services:

- *Predisposing factors*: are those factors that vary an individual's inclination to use services. These predisposing factors include demographic variables such as age and gender; social structure variables such as education, occupation, and ethnicity; and health beliefs about medical care, physicians, disease, and medication use.
- *Enabling factors*: those factors influencing the individual's ability to use services, thus they reflect the fact that an individual's ability to use services depends on individual family and community resources.
- *Need factors*: those factors related to the individual's belief in the seriousness of illness symptoms and the necessity of intervention. Need factors are separated into two categories, perceived need and evaluated need [5].

4. Factors That Influence Patient Behavior

4.1. Patient Factors

- a) **Age**: Older people tend to use health services more than younger people, they account for 34% of total pharmaceutical expenditures. Nearly 40% of the elderly have three or more chronic conditions, while about 1/3rd of the nonelderly have at least one chronic condition. Younger patients were more likely than older patients to know how their prescribed drugs worked, when it started working, common side effects, how to manage side effects, and how long their physician wanted them to take the medication [6].
- b) **Sex**: Women tend to use health services more than men. The self-reported use of Over The Counter (OTC) medications in the rural older population also shows that women take more OTC medications than men [7]. Sex also makes a difference in psychotropic medication use. Marked gender differences have been reported; women are affected twice as often as men [8]. In addition, women are twice as likely as men to report a positive family history of mood disorder, which is associated with a younger age of onset of depression [9]. Men receive more consultation from pharmacists. If considering the children, girls are encouraged to express their pain, whereas boys are encouraged to deny their pain and avoid feminine or sissy-like behaviors [10].
- c) **Socio-economic level**: Higher levels of both income and education are associated with lower rates of mortality [11]. Five themes emerged for patients who presented late for treatment: lack of knowledge, fear of hospitals, denial, living alone, and preexisting health concerns, found among stroke patients [12].
- d) **Ethnic Background**: *Jewish and Italian patients tended to have a more emotional response to pain*; they felt freer to discuss their pain, complain about it, groan and cry, and ask for relief. In contrast, patients

from other backgrounds tried to deny their pain and appear more stoic. The primary chronic health problem among Mexican Americans in the US is non-insulin-dependent diabetes mellitus [13].

e) **Communication Barriers, Financial Problems, And Cultural Barriers.**

- Latino patients are often very polite to doctors, so polite that rather than discuss their diabetes care, the patients nod their heads and agree with the doctor. Latino patients with limited English skills may be less likely to take prescribed diabetes medications than other diabetics in the U.S. The “nod of the head” in response to a physician’s instructions or comments may therefore represent a socially required gesture of respect, rather than understanding or agreement [14].
- Patients often do not believe that the medication supplies are free and therefore do not take the necessary diabetic supplies as often as needed.
- Other patients believe that receiving government assistance in medical supplies will decrease chances of US citizenship [15].
- For the families that do pay for medical supplies, a different problem arises. Expenses for a woman’s needs often are considered secondary to the good of her family, and therefore expenditures for diabetes medications and supplies are considered less important than other family necessities [16, 17].
- Finally, traditional folk remedies, such as aloe, cactus, and garlic, compete with the use of prescribed diet and medications, because patients (and possibly providers) are not aware that treatments can be combined [18].

f) **Emotional Factors:** Particular concern when the patient has been diagnosed with a terminal illness, an illness with a social stigma, or an illness that requires change in daily behavior. Factors of concern to patients include ~

- Uncertainty of what to expect with this new illness or symptom
- Dependency on providers to give the best treatment and on family to help with daily life
- Fear of change and death; pain and discomfort
- Lack of privacy in physical examinations
- Loss of identity as a healthy person
- Isolation from usual support systems such as coworkers, teammates, and friends; *conveying social support* also is important, because people are more likely to trust or respond positively to another person if they have an emotionally satisfying relationship with that person [19, 20].

4.2. **Drug Factors**

Drug regimens can be complex. The complexity of a drug regimen often is measured in the ~

- Total number of medications taken daily,
- Number of daily doses,
- Duration of treatment,
- The extent to which the regimen is tailored to daily routines,
- The side-effect profile [21].

Medications may require special behaviors, for example:

- Having to take a dose 1 hr. Before or 2 hr. after a meal
- Avoiding foods that are common in the diet
- Taking doses three or more times in a day,

- Refrigerator storage, or skill in administration.
- In addition, just learning the name of the drug prescribed, purpose of the drug, proper dose, when to begin taking it, frequency of dosing, and when to stop treatment is complex.
- The complexity of a therapeutic regimen may prevent patients from adhering completely. Complex regimens may produce information overload [22].
- Alternatively, medications requiring behaviors that are difficult to fit into regular daily activities are less likely to be taken as prescribed by a patient [23].

4.3. Environmental Factors

Patients given more autonomy and opportunities for self-determination tend to show greater health and morale improvements. The structural layout of many community pharmacies does not include an area for private consultation and dialog between the patient and the pharmacist. In addition to this lack of privacy, pharmacists often experience other environmental barriers to meaningful interaction with their patients, including

- Insufficient supportive personnel,
- A heavy workload and backlog,
- People waiting to present prescriptions or receive pharmacist assistance,
- Incoming phone calls and requests for information or help from coworkers,
- Interns, and other staff, and
- Inadequate computer technology, software, and
- Preparation for new consultation roles [24].

4.4. Provider Patient Interaction

Patients can play an important role in improving patient safety by becoming actively involved in their health care. Patients can act as 'safety buffers' during their care but the responsibility for their safety must remain with the health care

professionals [25]. Clinical practice often fails to optimize patient participation in decisions about serious illness. Patients prefer that the physician have most of the control over decisions [26, 27]. Patients use three main sources of information when making decisions about their illness and treatment:

- Their personal experience with the illness and various treatments;
- Information obtained from family, friends, and the larger culture;
- And their interaction with health professionals.

The patient-clinician relationship has both emotional and informational components – termed emotional care and cognitive care [28]. Emotional care includes mutual trust, empathy, respect, genuineness, acceptance and warmth [29]. Cognitive care includes information gathering, sharing medical information, patient education, and expectation management. Providers who adopt an autocratic approach assume a dominant or controlling role, speaking with an authoritarian tone and giving directions without seeking patient input. In contrast, providers who adopt a participatory approach collaborate with the patient to develop a mutually acceptable treatment plan, *providing decisional support* or guidance without ignoring patient views and demanding compliance with a certain therapeutic plan [30].

a) Provider Instruction on Patient Comprehension and Recall

Physicians and pharmacists continue to be the main sources of drug information and advice given to patients. Patients often receive information about the drug name and recommended dose and dosage frequency, but the majority of patients still receive no specific oral counseling about the purpose of therapy, how long to take their medication, side effects, other precautions, and when the medication will begin to work [31]. In fact, the quality of medication instruction by a provider is a better predictor of patient comprehension and recall than the patient's age and education. Limited

literacy has repeatedly been linked to problems comprehending health information, although the majority of studies to date have focused on reading various print health materials. Many older adults may have difficulty remembering verbal instructions conveyed during clinical encounters. Greater provider awareness of the impact of low health literacy on the recall of spoken instructions may guide providers to communicate more effectively and employ strategies to confirm patient understanding [32]. Research also has shown that there are substantial gains in patient comprehension and recall when providers use~

- Written reinforcement and visual aids, including printed leaflets or information sheets,
- Expanded prescription labels and stickers,
- Calibrated liquid measuring devices,
- And special containers or calendars that indicate exactly when each dose is to be taken.

It is estimated that close to half of the U.S. population has limited health literacy. Patients with Limited Health Literacy (LHL) are more likely to have poor health status, higher rates of hospitalization, and a nearly twofold higher mortality rate. They are also more likely to experience disparities in health and health care access, have lower rates of screening and preventive services, and obtain their care in the emergency department. Furthermore, patients with LHL are more likely to have poorer knowledge about their disease processes, medication regimens, and methods for managing their disease. LHL also has a negative effect on doctor-patient communication [33]. The difficulty and length of informational materials can interfere with the patient's ability to comprehend and recall advice. In general, patients have fewer difficulties if providers *simplify instructions* by avoiding medical jargon and using shorter words and sentences. In fact, the patients who received the difficult leaflet made nearly the same number of medication

errors as those who received no information [34, 35].

b) Provider Support on Patient Motivation and Evaluation of care

Being ill and undergoing treatment can involve

- A variety of stresses, practical problems, and
- Other concerns that adversely affect patients' evaluations of treatment and their motivation to perform difficult tasks such as changing an unhealthy life-style, taking multiple medications, tolerating adverse events, and maintaining a positive self-image and outlook

Patients also develop more positive attitudes and achieve better treatment outcomes when their caregivers make a systematic effort to *reinforce* the value of therapy. This reinforcement can take multiple forms, such as

- Giving feedback to patients about their conditions during follow-up medical and pharmacy visits,
- Encouraging patients to monitor their own conditions with special devices, or making home visits to increase family support and reinforcement. For example, experimental studies in hypertension management have documented substantial gains in patient adherence and clinical outcomes if patients receive regular blood pressure monitoring and feedback about their condition from a pharmacist or nurse [36, 37].

c) Provider Monitoring on Patient Feedback and Satisfaction

Surveys suggest that patients experience a wide variety of subjective and objective problems and concerns that contribute to nonadherence, dissatisfaction with care, and treatment dropout. These *barriers* to treatment adherence include ~

- Doubts about the physician's diagnosis or need for treatment,
- Misunderstandings about the regimen,
- Difficulties remembering each dose,

- Doubts about the effectiveness of the prescribed drug for their condition,
- Concerns about side effects and other bothersome features of a drug,
- And fears about the long-term effects of treatment or social stigma associated with certain conditions or treatments [38].

At the same time, physicians and pharmacists do not always ask patients about their medication concerns, beliefs, understandings, and behaviors. While these patient concerns can have detrimental effects on the patient's behavior, many patients are reluctant to complain or ask their providers about their medications. Providers who ask carefully designed, open-ended questions about patient concerns and adherence will be more likely to receive accurate patient feedback than those who wait for their patients to volunteer this information. community pharmacists should equip themselves with appropriate knowledge and competencies in order to tender efficient and outstanding pharmaceutical health care [39, 40].

5. Behavior: An East West Analogy

As reported by the WHO, patient adherence to long-term therapies is alarmingly low in both developed and developing countries [41]. Nearly 50% of antibiotics are purchased and consumed without any medical supervision in most parts of the world [42]. Evil spirits are blamed for epilepsy in most part of the world. Healthcare providers' understanding of their patients' healthcare beliefs, values, and preferences is an important feature of patient-centered care. The use of herbal medicinal products and supplements has increased tremendously over the past three decades with not less than 80% of people worldwide relying on them for some part of primary healthcare [43]. Medication nonadherence remains a substantial public health problem. Worldwide, between 25% and 50% of patients do not take their medications as recommended. In the USA, suboptimal adherence has been associated with 125,000 deaths, 10% of hospitalizations, and costs 300 billion USD annually [44]. The economic impact

was also in PubMed and Scopus in September 2017, which shows an increase in total healthcare cost (>80%), pharmacy costs (70%), inpatient and outpatient costs (50%), emergency department visit and medication costs (<30%), and hospitalization costs (<20%) [45]. Across South Asia, overall hypertension prevalence is estimated to be 27%. Prospective Urban Rural Epidemiology study has shown more than 50% are unaware of it and up to 80% of hypertensive patients have low adherence to medication [46]. Uncontrolled BP was found more than 50% in Bangladesh, 70% in Pakistan and almost 60% in Sri Lanka [47]. American Diabetes Association and the European Association for the Study of Diabetes guidelines in 10 developing countries from Africa (Egypt, South Africa), Middle East (Israel, Saudi Arabia, United Arab Emirates, Iran and Lebanon) and South Asia (Bangladesh, India and Pakistan) shows more than 25% and around 70% patients did not have any follow-up visit by a diabetologist or a GP, respectively. More than 35% patients did not receive any diabetes education [48].

6. Behavior: An East West Comparison

6.1. Health Expenditure and Utilization in Western Countries

- 25% of Hispanic patients report fear that insulin causes blindness. Patients need information that may not appear obvious to providers [49].
- In 2015, nearly \$3 trillion USD were spent on healthcare in the United States. 5% of the total population consumes 50% of all healthcare spending [50]. Just over half (54%) of healthcare funding in the United States comes from private sources, in contrast to just under one-third in Canada (30%) and Australia (32%) and under one-fifth in the United Kingdom (18%). Public sector spending per capita is more consistent across countries (Canada's expenditure is 5% greater than that of the United Kingdom and 20% greater than Australia's). During the year 2016, 91.2% of Americans had health

insurance coverage. Overuse has been best documented HICs [51].

- Australia has the highest number of MRI machines among OECD countries (13.4) per million people but has the lowest MRI exams (27.6) per 1000 people [52].

6.2. Misbeliefs, Superstitions and Poor Compliance in South Asian Countries

In Pakistan, 10% ethnic respondents rely on arm bracelet for pressure control and neem (*Azadirachta indica*) for diabetes measure; nearly 30% rely neem chicken pox and measles cure; 16% went to maulvis for treatment of jaundice; 20% of patients were made to smell shoes and 11% were taken to peers and faqirs (folk healers) [53]. “Coining”, a similar belief in China, Indonesia and many South Asian countries that rubbing coin along with the skin will bring bad blood to surface. Culturally in many parts of India, menstruation is still considered to be dirty and impure. A nearly 90% of women in India sometimes resort to using ashes, newspapers, dried leaves and husk sand to aid absorption [54]. The trend towards finding healers first and then doctors covers all fields of medicine in India and beyond. It cannot be wrong to show respect for cultural traditions and belief, but if pursued without heed to possible harms that arise from not seeking timely professional help the situation could change. Furthermore, the situation described is not unlike that which is commonly found in China, where the two different types of medicine (traditional and allopathic) generally work in parallel, and have done for many years. In Bangladesh, diarrhea and pneumonia cause death of more than ten lac children every year. Majority parents seek treatment from homeopaths, folk healers, retailers or non-qualified allopath [55]. A recent study shows that half of the antibiotics are sold without prescription in most of the South Indian countries, people prefer self-medication rather than visiting a doctor [56].

7. Pharmacists’ Role

7.1. Pharmacists Role in Improving Patient Behavior

1. Establish a relationship
2. Improve the patient and caregiver knowledge base
3. Utilizing digital health technology tools.
4. Providing incentives
5. Involve the caregiver [57], [58]

7.2. Importance of Patient Counseling

Patient counseling is necessary to reduce medication errors and improve patient healthcare. This leads to several potential benefits:

- Improved therapeutic outcomes and decreased adverse effects
- Improved patient adherence to the treatment plan
- Decreased medication errors and misuse
- Enhanced patient self-management by involving the patient in designing the therapeutic plan

Potential for decreased health care costs due to appropriate use of medications and prevention of adverse events. The pharmacist also benefits in this process. Potential benefits to the pharmacist in this process include:

- Enhanced professional status in the view of patients and other health care providers
- Establishment of an essential component of patient care that cannot be replaced by technicians or automation
- Enhanced job satisfaction through improving patient outcomes
- A value-added service to offer patients
- Revenue generation through payment for counseling services– limited at present but growing

In short, it ensures positive outcomes on the management of disease, including improved drug compliance, better treatment endpoints, and patient satisfaction [69].

7.3. Benefits of Patient Compliance

The improvement of compliance will result in a situation in which all parties benefit. Most

importantly patients benefit from the enhancement of the efficacy and safety of their drug therapy. Pharmacists benefit because there is an increased recognition and respect for the value of the advice and service that they provide. Pharmaceutical manufacturers benefit from the favorable recognition that accompanies the effective and safe use of their drugs as well as from the increased sales resulting from the larger number of prescriptions being dispensed. Finally, society and the health care system benefit as a result of fewer problems associated with noncompliance. Although an increase in compliance will result in more prescriptions being dispensed and a higher level of expenditures for prescription medications, this increase in costs will be more than offset by a reduction in costs (e.g., physician visits, hospitalizations) attributable to problems due to noncompliance [60], [61].

7.4. Conflict Resolution and Problem Solving

Because providers and patients are likely to have different viewpoints and agendas, some interpersonal conflict or disagreement is inevitable during their interaction. This conflict is especially likely after patients have gained more experience with their illness and recommended treatments. For example, providers may want patients to comply as fully and rapidly as possible with their ideal treatment plan, whereas patients may prefer a slower or less aggressive approach or even request alternative therapies that providers would consider ineffective, inappropriate, or unnecessary. Providers also may consider certain side effects or drug-taking problems to be clinically insignificant or trivial, while patients consider the same side effects or problems to be intolerable. As a result, providers often are confronted with both explicit and subtle forms of negative feedback from their patients (eg, complaints about the drug or dosage schedule, admissions of nonadherence, reported difficulties administering or paying for medication, expressions of fear and uncertainty about drug efficacy or safety). In contrast, the participatory/collaborative approach involves

acknowledging the legitimacy of patient concerns, assessing patient concerns in a more thorough and respectful manner, tailoring or adjusting drug regimens to fit patient routines and preferences, and negotiating mutually acceptable solutions. Patient adherence is significantly improved if the dosage schedule has been tailored to the patient's daily routine, if the patient is allowed to change the regimen within a preapproved protocol, and if the patient identifies the areas in which he or she would like assistance [62], [63].

7.5. The Health Collaboration Model

Health Collaboration Model highlights the central role of patient feedback and collaborative problem solving in health care. Each box represents a different set of provider behaviors or background factors affecting the collaboration process; whereas, each circle represents a different set of patient cognitions, beliefs, behaviors, or clinical outcomes affected by collaboration and background factors [64]. Health Collaboration Model highlights the central role of patient feedback and collaborative problem solving in health care. The diagram is best understood by beginning at the left side of the diagram and following each arrow in numerical order. Each box represents a different set of provider behaviors or background factors affecting the collaboration process; whereas, each circle represents a different set of patient cognitions, beliefs, behaviors, or clinical outcomes affected by collaboration and background factors [65].

The top part of the model (arrows 1–5) emphasizes the various factors that can impact the patient's initial reactions after receiving a new prescription or other new regimen; whereas, the bottom part of the model (arrows 6–13) emphasizes the various factors that can impact the patient's behavior after he/she has experienced the drug and barriers to adherence. First, we see that patient, provider, drug, and environmental factors (arrow 1) can impact both provider and patient behavior during the initial stages of collaboration and treatment, including the quality of provider instruction and support, patient comprehension/ recall and motivation, and initial

patient adherence and barriers to adherence. During the initial stages of collaboration, the provider plays a critical role in facilitating and verifying patient comprehension/recall of the regimen (arrow 2). The provider also plays a critical role in facilitating and verifying the patient's initial motivation and satisfaction with the regimen (arrow 3). It is not

enough to give drug information. Rather, the provider must assess the individual's initial understandings and beliefs and make adjustments as necessary. In some cases, he/she will need to provide additional instruction or reinforcement to make sure the patient will understand and remember the dosage schedule.




Figure 1. Health collaboration model [66].

In other cases, he/she will need to provide additional informational, social, or technical support to address the individual's initial doubts or concerns about the drug and its short-term or long-term effects. If effective collaboration occurs at this stage, then the patient will have greater comprehension/recall and greater motivation which, in turn, lead to greater initial adherence and fewer barriers to adherence (arrows 4–5). The bottom part of the model draws attention to the fact that patients actively monitor their reactions to drug therapy and experience a variety of barriers to adherence after initiating therapy. They may find it difficult to remember each dose or simply question whether the drug is still needed or working for them. They also may experience unwanted effects or have concerns about high drug costs, possible dependence, or unknown long-term

effects. Since these patient-perceived barriers seriously undermine the patient's willingness and ability to continue therapy as prescribed, patient feedback plays a central role in follow-up visits with the provider (as shown in Fig 1). Soliciting accurate patient feedback is a complex process that depends on: the patient's personal experience with the drug (arrow 6); background characteristics and expectations of the patient and provider, characteristics of the drug and environment, and past patient-provider interactions (arrow 7); and quality of provider monitoring. Providers who regularly ask carefully designed, open-ended questions in a supportive, non-accusatory manner can be very successful in soliciting negative patient feedback even among patients who initially appear reluctant or hesitant to share their personal doubts, difficulties, and concerns. The provider

who is able to solicit accurate patient feedback is then able to resolve patient-specific problems and provide appropriate reinforcement as necessary (arrows 9–10). This type of two-way communication and collaborative problem-solving leads to greater patient satisfaction and adherence (arrow 11) and enhanced treatment outcomes (arrow 12). The final arrow (14) illustrates the importance of past interactions and treatment experiences in establishing and maintaining a trusting relationship that is the cornerstone of effective health and pharmaceutical care.

Health Collaboration Model can play an important role in pharmacy practice and research.

- It enables pharmacy practitioners and researchers to organize large amounts of information that would otherwise be confusing or difficult to interpret and use.
- It enables pharmacists to identify potential connections and implications that are not obvious when examining results from a single study or set of observations.
- It can be used as a stimulus and guide for further discussion, evaluation, and practice development.
- It helps us see that the patient's behavior depends more upon the patient's beliefs, feelings, and interactions than on the patient's medical diagnosis or severity of illness.
- It also helps us see pharmacists who have a good understanding of patient behavior can have a positive impact on treatment outcomes by providing quality instruction, support, monitoring, and collaborative problem-solving and reinforcement [67], [68].

8. Conclusion

The comparative study shows different perspective of patient behavior in modern world and in Bangladesh. Reasons are found to cultural and economic differences. Pharmacists and other healthcare providers also have

different strategies to promote patient behavior change, spent times with patients. High patient activity change support system should validate their efficiency and professionalism. But reality is that no graduate pharmacist is working in retail pharmacies or Government hospitals of Bangladesh except very few tertiary private hospitals. To attain the ideal healthcare service doctor, pharmacist, nurse and other healthcare professionals must work together. It is noteworthy to mention that any interruption in the team work will disrupt the whole health system and patient care service will never be achieved. At the same time, physicians and pharmacists do not always ask patients about their medication concerns, beliefs, understandings, and behaviors. While these patient concerns can have detrimental effects on the patient's behavior, many patients are reluctant to complain or ask their providers about their medications. Providers who ask carefully designed, open-ended questions about patient concerns and adherence will be more likely to receive accurate patient feedback than those who wait for their patients to volunteer this information.

Acknowledgement

It's a great gratitude and honor to be a part of healthcare research and pharmacy education. Pharmacists and health providers of all disciplines that I have conducted was very much helpful in discussing healthcare situations in Bangladesh and abroad, providing books, journals, newsletters and precious time. The greatest help was from my students who paid interest in my topic as class lecture and encouraged to write such article comprising patient communication and behavior management. Despite a great scarcity of funding this purpose from any authority, the experience was good enough to carry on research.

Compliance with The Ethical Issues

•Ethics approval and consent to participate

Animal and Human experiment: N/A

Human Data Submission Approval: N/A

•Consent for publication

Consent to publish Individual Person's data: N/A

•Availability of data and materials

Data sharing: Data will be provided on request.

•Competing interests

The author declares that he has no competing interests.

•Funding

Funding from individual/Organization: No fund received from any individual/organization.

•Authors' contributions

The individual contributions of authors: N/A

References

1. Manmohan T, Sreenivas G, Sastry VV, Sudha ER and others. DRUG COMPLIANCE AND ADHERENCE TO TREATMENT Journal of Evolution of Medical and Dental Sciences/Volume1/ Issue3/July-Sept 2012 Page 142-159
2. Sarieh P, Afsaneh R, Parisa B, Samad EJG, Soroor P, Forough R. Evolutionary concept analysis of health seeking behavior in nursing: a systematic review BMC Health Serv Res. 2015; 15: 523. doi: [10.1186/s12913-015-1181-9] PMID: 26613729
3. Irwin MR. Why People Use Health Services Milbank Q. 2005 Dec; 83(4): 10.1111/j.1468-0009.2005.00425.x. doi: [10.1111/j.1468-0009.2005.00425.x] PMID: PMC2690262
4. Stanislave VK, Sidney C. Health Behavior, Illness Behavior, and Sick-Role Behavior II. Sick-Role Behavior Archives of Environmental Health: An International Journal Volume 12, 1966 - Issue 4 Pages 531-541 | Received 10 Nov 1965, Accepted 17 Nov 1965, Published online: 30 Apr 2013
<https://doi.org/10.1080/00039896.1966.10664421>
5. Han KK, Munjae L. Factors associated with health services utilization between the years 2010 and 2012 in Korea: using Andersen's Behavioral model Osong Public Health Res Perspect. 2016 Feb; 7(1): 18–25. doi: [10.1016/j.phrp.2015.11.007] PMID: 26981338
6. Mueller C, Schur C, O'Connell J. Prescription drug spending: the impact of age and chronic disease status. Am J Public Health. 1997 October; 87(10): 1626–1629. PMID: 9357343
7. Jennifer P, Ineke N. Robert C. Marie DM. BMC Womens Health. 2004; 4(Suppl 1): S29. doi: [10.1186/1472-6874-4-S1-S29] PMID: 15345092
8. Kessler RC (2003). Epidemiology of women and depression. Journal of Affective Disorders 74, 5–13. [https://doi.org/10.1016/S0165-0327\(02\)00426-3](https://doi.org/10.1016/S0165-0327(02)00426-3)
9. Nierenberg AA, Trivedi MH, Fava M, Biggs MM, Shores-Wilson K, Wisniewski SR, Balasubramani GK, Rush AJ. (2007). Family history of mood disorder and characteristics of major depressive disorder: A STAR (*) D (sequenced treatment alternatives to relieve depression) study. Journal of Psychiatric Research 41, 214–221. <https://doi.org/10.1016/j.jpsychires.2006.02.005>
10. Brannon R, David DS. The male sex role: Our culture's blueprint of manhood and what it's done for us lately. The forty-nine percent majority: The male sex role. Reading, MA: Addison-Wesley.
11. Robert AH, Elaine MH. The Effect of Educational Attainment on Adult Mortality in the United States. Popul Bull. 2013 Jun; 68(1): 1–16. PMID: 25995521
12. Kitko L, Hupcey JE, Factors that influence health-seeking behaviors of patients experiencing acute stroke. J Neurosci Nurs. 2008 Dec;40(6):333-40.
13. Gary BR. CHAPTER 6. Ethnocultural Variations in the Experience of Pain. 2004, English, Book, Illustrated edition: Pain: psychological perspectives / edited by Thomas Hadjistavropoulos, Kenneth D. Craig. Bookmark: <https://trove.nla.gov.au/version/28877025> Published Mahwah, N.J.; London : Lawrence Erlbaum, 2004.
14. Pachter, LM. Cultural issues in pediatric care. in: Nelson's textbook of pediatrics.: Springer, New York; 1992
15. Alexander NO, Hector PR, Arturo VB. Policy Dilemmas in Latino Health Care and Implementation of the Affordable Care Act Annu Rev Public Health. 2015 Mar 18; 36: 525–544. doi: [10.1146/annurev-publhealth-031914-122421] PMID: 25581154
16. Martz DM, Handley KB, Eisler RM. The relationship between feminine gender role stress, body image, and eating disorders. Psychology of Women Quarterly. 1995; 19: 493-508
17. Muhammad AS, Mannan FK, Thomas EC. Gender Differences in Living with Diabetes Mellitus Mater Sociomed. 2013; 25(2): 140–142. doi: [10.5455/msm.2013.25.140-142] PMID: 24082841
18. Hannah N, Dara HR and others. Complementary and Alternative Medicine (CAM) Use among Non-Hispanic White, Mexican American, and Vietnamese American Patients with Type 2

- Diabetes J Health Care Poor Underserved. 2014 Nov; 25(4): 1941–1955. doi: [10.1353/hpu.2014.0178] PMID: 25418251
19. Jane T, Brian K. Emotional dimensions of chronic disease. *West J Med.* 2000 Feb; 172(2): 124–128. PMID: 10693376
 20. Haifeng L, Yang J, Tianyong C. The Roles of Different Sources of Social Support on Emotional Well-Being among Chinese Elderly *PLOS* Published: March 3, 2014 <https://doi.org/10.1371/journal.pone.0090051>
 21. Beena J, Jimmy J. Patient Medication Adherence: Measures in Daily Practice *Oman Medical Journal* (2011) Vol. 26, No. 3: 155-159 *Oman Medical Journal* (2011) Vol. 26 URL: <https://www.ncbi.nlm.nih.gov/pmc/articles/PMC3191684/>
 22. Ian MK, Siqin Y. Adherence to Cardiovascular Medications: Lessons Learned and Future Directions. *Prog Cardiovasc Dis.* 2013 May-Jun; 55(6): 590–600. doi: [10.1016/j.pcad.2013.02.001] PMID: 23621969
 23. Hayden BB, Bradi BG. Medication Adherence: A Call for Action. *Am Heart J.* 2011 Sep; 162(3): 412–424. doi: [10.1016/j.ahj.2011.06.007] PMID: 21884856
 24. Laetitia HH, Lynne E, Pascale NCT, Catherine G. Utilization of community pharmacy space to enhance privacy: a qualitative study *Health Expect.* 2016 Oct; 19(5): 1098–1110. doi: [10.1111/hex.12401] PMID: 26332335
 25. Davis RE, Jacklin R, Sevdalis N, Vincent CA. Patient involvement in patient safety: what factors influence patient participation and engagement? *Health Expectations*, 10, pp.259–267
 26. Ende JL. Preferences for autonomy when patients are physicians. *J Gen Intern Med.* 1990;5506-509
 27. Bradley J, Hamilton N. Patient preferences for control in medical decision making: a scenario-based approach. *Fam Med.* 1996;28496- 501
 28. Blasi DZ, Harkness E, Ernst E, Georgiu A, Kleijnen J (2001) Influence of context effects on health outcomes: A systematic review. *Lancet* 357: 757–762
 29. Ong LML, de Haes JCJM, Hoos AM, Lammes FB (1995) Doctor-patient communication: A review of the literature. *Social Science & Medicine* 40: 903–918.
 30. National Academies Press (US): 2006 Chapter 5. Individuals and Families: Models and Interventions. *Health and Behavior: The Interplay of Biological, Behavioral, and Societal Influences.* Institute of Medicine (US) Committee on Health and Behavior: Research, Practice, and Policy, Washington (DC)
 31. Zeng-Treitler Q, Kim H, Hunter M. Improving patient comprehension and recall of discharge instructions by supplementing free texts with pictographs. *AMIA Annu Symp Proc.* 2008; 2008: 849–853. Published online 2008. PMID: 18999109
 32. Danielle mc, Katherine RW. What Did the Doctor Say? Health Literacy and Recall of Medical Instructions. *Med Care.* 2012 Apr; 50(4): 277–282. doi: [10.1097/MLR.0b013e318241e8e1] PMID: 22411440
 33. Rebecca LS, Dean S. Interventions to Improve Care for Patients with Limited Health Literacy. *J Clin Outcomes Manag.* 2009 Jan 1; 16(1): 20–29. PMID: 20046798
 34. Sameer B, Sanjeev S. Assessing Readability of Patient Education Materials: Current Role in Orthopaedics. *Clin Orthop Relat Res.* 2010 Oct; 468(10): 2572–2580. doi: [10.1007/s11999-010-1380-y] PMID: 20496023
 35. National Academies Press (US): 2004. Chapter 6. Health Systems. *Health Literacy: A Prescription to End Confusion.* Institute of Medicine (US) Committee on Health Literacy; Nielsen-Bohlman L, Panzer AM, Kindig DA, editors. Washington (DC)
 36. Derek CS, Johnson G and others. Views of pharmacist prescribers, doctors and patients on pharmacist prescribing implementation. *International Journal of Pharmacy Practice* 2009; 17: 89–94
 37. Manabu A, Akiko M and others. Establishing a Pharmacy-Based Patient Registry System: A Pilot Study for Evaluating Pharmacist Intervention for Patients with Long-Term Medication Use. *Pharmacy (Basel).* 2018 Mar; 6(1): 12. doi: [10.3390/pharmacy6010012] PMID: 29370145
 38. Elizabeth JU. Development of Models to Predict Medication Non-adherence Based On a New Typology. Thesis and dissertations University of Iowa. Iowa Research Online.
 39. Deborah VK, Stephanie Y and others. Patient attitudes regarding the role of the pharmacist and interest in expanded pharmacist services. *Can Pharm J (Ott).* 2014 Jul; 147(4): 239–247. doi: [10.1177/1715163514535731] PMID: 25360150
 40. Mohamed NA. Patients' perception, views and satisfaction with pharmacists' role as health care provider in community pharmacy setting at Riyadh, Saudi Arabia. *Saudi Pharm J.* 2012 Oct; 20(4): 323–330. doi: [10.1016/j.jsps.2012.05.007] PMID: 23960807
 41. Anghel LA, Farcaş AM, Oprean RN. Medication adherence and persistence in patients with autoimmune rheumatic diseases: a narrative review. *Patient Prefer Adherence.* 2018;12:1151-

1166. Published 2018 Jul 3. doi:10.2147/PPA.S165101
42. Nepal G, Bhatta S. Self-medication with Antibiotics in WHO Southeast Asian Region: A Systematic Review. *Cureus*. 2018;10(4):e2428. Published 2018 Apr 5. doi:10.7759/cureus.2428
 43. The growing use of herbal medicines: issues relating to adverse reactions and challenges in monitoring safety. *Front Pharmacol*. 2014;4:177. Published 2014 Jan 10. doi:10.3389/fphar.2013.00177
 44. Zullig LL, Blalock DV, Dougherty S, et al. The new landscape of medication adherence improvement: where population health science meets precision medicine. *Patient Prefer Adherence*. 2018;12:1225-1230. Published 2018 Jul 13. doi:10.2147/PPA.S165404
 45. Cutler RL, Fernandez-Llimos F, Frommer M, Benrimoj C, Garcia-Cardenas V. Economic impact of medication non-adherence by disease groups: a systematic review. *BMJ Open*. 2018;8(1):e016982. Published 2018 Jan 21. doi:10.1136/bmjopen-2017-016982
 46. Naheed A, Haldane V, Jafar TH, Chakma N, Legido-Quigley H. Patient pathways and perceptions of hypertension treatment, management, and control in rural Bangladesh: a qualitative study. *Patient Prefer Adherence*. 2018;12:1437-1449. Published 2018 Aug 14. doi:10.2147/PPA.S163385
 47. Jafar TH, Gandhi M, Jehan I, Naheed A, de Silva HA and others. COBRA-BPS Study Group; Determinants of Uncontrolled Hypertension in Rural Communities in South Asia—Bangladesh, Pakistan, and Sri Lanka, *American Journal of Hypertension*, Volume 31, Issue 11, 15 October 2018, Pages 1205–1214, <https://doi.org/10.1093/ajh/hpy071>
 48. Al Mansari A, Obeid Y, Islam N, et al. GOAL study: clinical and non-clinical predictive factors for achieving glycemic control in people with type 2 diabetes in real clinical practice. *BMJ Open Diabetes Res Care*. 2018;6(1):e000519. Published 2018 Jul 10. doi:10.1136/bmjdr-2018-000519
 49. Jie H, Karen A. The meaning of insulin to Hispanic immigrants with type 2 diabetes and their families. *Diabetes Educ*. 2012 Mar-Apr; 38(2): 263–270. doi: [10.1177/0145721712437559] PMID: 22357711
 50. Clewley D, Rhon D, Flynn T, Koppenhaver S, Cook C. Health seeking behavior as a predictor of healthcare utilization in a population of patients with spinal pain. *PLoS One*. 2018;13(8):e0201348. Published 2018 Aug 1. doi:10.1371/journal.pone.0201348
 51. Brownlee S, Chalkidou K, Doust J, et al. Evidence for overuse of medical services around the world. *Lancet*. 2017;390(10090):156-168.
 52. Dixit SK, Sambasivan M. A review of the Australian healthcare system: A policy perspective. *SAGE Open Med*. 2018;6:2050312118769211. Published 2018 Apr 12. doi:10.1177/2050312118769211
 53. Bukhari A, Pardhan A and others. Superstitions regarding Health Problems in different EthnicGroups in Karachi. URL: <http://www.jpma.org.pk/PdfDownload/2369.pdf>
 54. Garg S, Anand T. Menstruation related myths in India: strategies for combating it. *J Family Med Prim Care*. 2015;4(2):184-6.
 55. Lana V, Shahnawaz A, Farzana F and others. Self-care practices and barriers to compliance among patients with diabetes in a community in rural Bangladesh *Int J Diabetes Dev Ctries* DOI 10.1007/s13410-015-0460-7
 56. Kumar BS, Anamika S., Naz HH, and Riaz U. Irrational Use of Antibiotics and Antibiotic Resistance in Southern Rural Bangladesh: Perspectives from Both the Physicians and Patients *Annual Research & Review in Biology* 4(9): 1421-1430, 2014
 57. Paul R. Role of community pharmacists in patients' self-care and self-medication. *Integr Pharm Res Pract*. 2015; 4: 57–65. doi: [10.2147/IPRP.S70403] PMID: 29354520
 58. Jolly F, Suja A. Clinical pharmacists: Bridging the gap between patients and physicians. *Saudi Pharm J*. 2014 Dec; 22(6): 600–602. doi: [10.1016/j.jsps.2014.02.011] PMID: 25561874
 59. Sanii Y, Torkamandi H. Role of pharmacist counseling in pharmacotherapy quality improvement. *J Res Pharm Pract*. 2016 Apr-Jun; 5(2): 132–137. doi: [10.4103/2279-042X.179580] PMID: 27162808
 60. Mohiuddin AK. Patient Compliance: An Untold Story or a Fairy Tale? *Pharmaceut Reg Affairs* 2018, 7:2 DOI: 10.4172/2167-7689.1000207 ISSN: 2167-7689
 61. Christensen-Szalanski JJ, Northcraft GB. Patient compliance behavior: the effects of time on patients' values of treatment regimens. *Soc Sci Med*. 1985;21(3):263-73. PMID 3929392.
 62. Michael AER. Conflict in the health care workplace *Proc (Bayl Univ Med Cent)*. 2001 Apr; 14(2): 138–139. PMID: 16369603
 63. Rose-Lima VK, Reginald D and others. Conflicts between healthcare professionals and families of a multi-ethnic patient population during critical care: an ethnographic study. *Crit Care*. 2015; 19: 441. doi: [10.1186/s13054-015-1158-4] PMID: 26694072

64. Mays GP, Halverson PK, Kaluzny AD. Collaboration to improve community health: trends and alternative models. *Jt Comm J Qual Improv.* 1998 Oct;24(10):518-40. PMID 9801951.
65. National Academies Press (US): 2013. Chapter 7. Engaging Patients, Families, and Communities. *Best Care at Lower Cost: The Path to Continuously Learning Health Care in America.* Committee on the Learning Health Care System in America; Institute of Medicine; Smith M, Saunders R, Stuckhardt L, et al., editors. Washington (DC). May 10, 2013.
66. Pharmaceutical Press Website. *The Health Collaboration Model: A Tool for Analyzing and Improving Patient Care.*
67. CSHP. *Pharmacy Practice in Hospitals and Other Collaborative Healthcare Settings: Position Statements (2016).* Official Publication, Position Statement of Canadian Society of Hospital Pharmacists (CSHP), Ottawa, Ontario. 2016. © Canadian Society of Hospital Pharmacists.
68. Benjamin P, Robert JW. Enhancing Pharmacy Practice Models Through Pharmacists' Privileging. *Hosp Pharm.* 2013 Feb; 48(2): 160–165. doi: [10.1310/hpj4802-160] PMID: 24421454

