AJFRR 2018,1:4 **Research Article** 



## American Journal of Forensic Research and Reviews (DOI:10.28933/AJFRR)



# To identify a prevalence of suicide attempt and treatment, clinical, sociodemographic and epidemiological profile of adolescents

Andrade R.M.A1, Vasconcelos R.P2, Konze V.S2, Moraes S.R.T4, Souza M.F.M5, Ramos R.C.F6

1.2.3.4Estudantes do Curso de Medicina – Unicap: 5.6Docentes do Curso de Medicina – Unicap

### **ABSTRACT**

Introduction: It is estimated that for each suicide, there are 10 to \*Correspondence to Author: 20 attempts that often leave a high sequelae rate in the health Andrade R.M.A and physical incapacity of the individuals who cannot achieve it. Estudantes do Curso de Medicina Currently, there is a greater concern regarding some age groups, - Unicap such as adolescence, the risk classification in about one third of the countries. Objectives: To identify a prevalence of suicide attempt and treatment, clinical, sociodemographic and epidemi- How to cite this article: ological profile of adolescents involved in this picture. Methodol- Andrade R.M.A, Vasconcelos R.P. ogy: A cross-sectional, descriptive and retrospective study about Konze V.S, Moraes S.R.T, Souza suicide attempt in adolescents from the metropolitan region of M.F.M, Ramos R.C.F.To identify a Recife and attended at an outpatient clinic specialized in He- prevalence of suicide attempt and biatrics in the city of Recife, Pernambuco, between May 2016 treatment, clinical, sociodemoand August 2017. Results and Discussion: 8% of the patients at- graphic and epidemiological profile tempted suicide, these being more females. As motivation were of adolescents, American Journal reported: depression, bullying/stress, dissatisfaction with the of Forensic Research and Reviews, body and parents' anger were reported. The majority reported 2018,1:4. religious practice and was embedded in youth groups. Conclusion: Despite the restricted sampling, a prevalence of attempted suicide among adolescents was published in other countries. A family situation is still worrying if there are internal disorders. The insertion of the young person into daily activities is essential for the reduction of suicide attempts.

**Keywords:** Adolescent, Prevalence, Suicide attempt

