



Global Journal of Energy and Environment (DOI:10.28933/GJEE)



DEPLETION IN THE BIOMASS OF FLORAL WASTE-CATTLE DUNG MIXTURES DURING VERMICOMPOSTING

*Praveesh Bhati¹ Ph.D., Shobha Shouche² Ph.D. and Sudhir Kumar Jain³ Ph.D.

¹Ex Faculty, Govt. Madhav Science College, Dewas road Ujjain (M.P.) 456010, India; ² Assistant Professor, Govt. Madhav Science College, Affiliated to Vikram University Dewas road Ujjain (M.P.) 456010, India; ³ Reader, SOS in Microbiology, Affiliated to Vikram University Dewas road Ujjain (M.P.) 456010, India

ABSTRACT

Vermicomposting of different proportion of floral waste (FW)-cattle dung (CD) mixtures were carried out under aerobic condition in a plastic bins to find the rate of depletion of biomass. The initial substrate depth of five different compositions viz. 100% FW, 75% FW, 50% FW, 25% FW and 100% CD were 25cm, 16 cm, 13 cm, 11 cm and 8 cm respectively. The rate of biomass reduction was high in 100% FW which was decreased with increasing the quantity of cattle dung. The time for substrate stabilization during vermicomposting was also found to be lesser in summer as compared to rainy season. The substrate depth variation with respect to time followed a declined curve.

Keywords: Biomass reduction, floral waste, cattle dung, vermicomposting, earthworm.

*Correspondence to Author:

Praveesh Bhati

Ex Faculty, Govt. Madhav Science College, Dewas road Ujjain (M.P.) 456010, India:

How to cite this article:

Praveesh Bhati Shobha Shouche and Sudhir Kumar Jain. Natural resource modelling: accounting for gillnet size selectivity in dynamic deterrence model. Global Journal of Energy and Environment, 2019,1:4.

 eSciPub
eSciPub LLC, Houston, TX USA.
Website: <http://escipub.com/>

Introduction:

Vermicomposting is a biotechnological process in which organic solid waste is subjected to breakdown in a moist, warm and aerated environment (Finstein and Morris, 1975; Haug, 1993; Narayan, 1993; Miller, 1996). Vermicomposting is an imperative aspect, as it transforms organic waste to prosperity product by using inexpensive eco-friendly opportunity with action of earthworm and microorganisms (Mall *et.al.*2005). Many types of solid waste, such as food, paper, wastewater sludge, and garden wastes, are acquiescent to biodegradation as they possess major amounts of heterogeneous organic substrates, including sugars (as sucrose), fats, proteins, hemicelluloses, celluloses, and lignins (Gray *et. al.*, 1971a; Rhyner *et. al.*, 1995; Eklind *et. al.*, 1997). When microorganisms *viz.* bacteria, fungi, and actinomycetes decompose organic materials in the presence of atmospheric oxygen, generates metabolic heat, carbon dioxide, NH₃ and water (Ryckeboer *et. al.*, 2003). Nitrogen and Carbon of the manure could be lost in the form of gas. (Tiquia and Tam 2000a). It was also found that during decomposition process, heat generated that decapitate in the form of water lost as a result biomass also reduced (Vogt, *et. al.*, 1986). It has been reported that about 40% volume of waste materials reduced from their initial value due to composting (Banu *et. al.*, 2008). Ndegwa and Thompson (2001) also suggested that during the process of composting and vermicomposting, biosolid waste reduced approximately 45% by their original weight. It is necessary to evaluated the ratio of initial volume of floral waste and finished good so that we can calculate the conversion ratio of solid organic waste.

The purpose of this paper is to describe loss of biomass of organic waste during vermicomposting and provide information about processes and data useful to predict the amount of organic waste taken into the bed for vermicomposting.

Materials and methods:

Collection of composting wastes:

Vermicomposting process was carried out of the cattle dung and floral waste. The cattle dung (CD) was collected locally from randomly selected cattle houses of municipal territory of the study area, Ujjain. The cattle dung was collected in large-sized rectangular plastic pot containers and was brought to vermicomposting unit, Government M.V.M., Ujjain. *Eudrilus eugeniae* and *Eisenia foetida* were equally used in this study, obtained from vermiculture centre of Govt. Madhav Science College, Ujjain (M.P.). The floral wastes (FW) was used in this experiment, collected from Shri Mahakaleshwer and Iskon temple. The non-recyclable items were separated through hand sorting from collected floral waste (Suthar and Singh, 2008).

Preparations of vermicomposting materials:

Collected floral wastes were chopped into small pieces. The chopped wastes were mixed with cattle dung in different ratio (Table no.1). Each composition was prepared in two sets. Composting bins were kept in bricks room. The windrow compost method was used in which it was not covered and ventilation was not provided with pipes. Bins were sprinkled with distilled water after turning it upside down to maintain high moisture content. Twenty earthworms (*E. eugeniae* and *E. foetida*) were added in each composting bins (Singh, *et.al.*2004). During process period, biomass reduction was measured.

Measurement of biomass reduction:

Reduction in the biomass of floral composting mixture was measured by plastic scale. This was done after interval of 24 hours throughout the complete process. Before the measurement, scale was disinfect by 70% alcohol and allowed to air dry. Composting materials were spread evenly in composting bins. Before taking the depth of composting materials, bins were jerk on the surface two times so that gap between floral wastes was



Photograph no.1: Measurement of biomass reduction of composting wastes.

Table -1: Representative mixture of floral waste –cattle dung in different compositions.

S.No.	Compositions	Floral waste	Cattle dung	Total volume
1.	100% Floral waste	2.0kg	0.0 kg	2.0 kg
2.	75% Floral waste	1.5kg	0.5 kg	2.0 kg
3.	50% Floral waste	1.0kg	1.0 kg	2.0 kg
4.	25% Floral waste	0.5kg	1.5 kg	2.0 kg
5.	100% Cattle dung	0.0kg	2.0 kg	2.0 kg




Figure 1 Depletion of biomass of floral waste-cattle dung mixtures in Rainy season.




Figure-II: Depletion of biomass of floral waste-cattle dung mixtures in Winter season.




Fig.-3: Depletion of biomass of floral waste-cattle dung mixtures in Summer season

Note: Series-1: 100 % floral wastes

Series-2: 75% floral waste-25% cattle dung

Series-3: 50 % floral waste – 50% cattle dung

Series-4: 25 % floral waste – 75% cattle dung

Series-5: 100 % cattle dung

filled with adjacent composting materials. Scale was kept vertically on the composting materials in all four sides of bin. From the surface of composting materials, length was measured to the edge of plastic bin. Mark was put at measuring site. After taken the four values, their mean was considered (Singh, *et. al.*, 2004).

Results and Discussions:

Based on the observation carried out for a period of 45 days in three seasons viz. rainy, winter and summer, the results of the reduction in the substrate column depth with respect to the time during the course of vermicomposting are presented in figures. The initial substrate depth of five different compositions viz. 100% FW, 75% FW, 50% FW, 25% FW and 100% CD were 25cm, 16 cm, 13 cm, 11 cm and 8 cm respectively. All the initial values were considered as control and any changed in the level of biomass of composting bins was determined as reduction level. Result obtained shows that in three seasons, among the five different composting mixtures, highest biomass was depleted in 100% FW which was up to 20 cm³ while lowest depletion was seen in 100% CD that was only 2 cm³. In the rainy season, (Figure I) biomass of vermicompost was decreased rapidly up to first twelve days of composting but soon after rate of depletion reduced. In all mixtures, depletion of biomass set at constant value (5 cm³) after 38th day of vermicomposting. In winter season (Figure II) the rate of biomass reduction was slowest as compared to rainy seasons. Initial 10 days biomass of floral –cattle dung waste depleted rapidly but after it became slower. After 35th day, biomass reductions of all mixtures were set around 5cm³. In summer season (Figure III), the rate of biomass reduction was highest. First nine day of vermicomposting, biomass was reduced rapidly but later it was set at constant value at 21th day. It was also noticed that in summer season, reduction rate of biomass highest and in rainy season, it was lowest.

The maximum depth reduction was took place in the first twelve days. The depth reduction rate subsequently slowed down during the 20 to 30 days of vermicomposting. Afterwards, (i.e., after about 30 days), almost no depth reduction was observed. It has been found that highest biomass reduced in 100% floral waste while it was minimum in 100 % Cattle dung. It was also revealed that the rate of biomass reduction affected by season, temperature, moisture content, nature of biomass and quantity of cattle dung. These reductions in the bulk density may be due to the high initial moisture content of the raw waste that was released in the form of water vapor. Along with this complex organic matter also turn down in to simple form therefore their density was reduced. Norbu (2002) also carried out experiment on municipal solid waste and found that, biomass of waste reduced rapidly in first two week then it was set as constant level. Singh, *et. al.* (2004) were also concluded that the maximum depth reduction takes place in the first seven days and later it was established after 25 days. Similar observation was also recorded by Moqsud (2010) who stated that composting was completed in about 35-40 days and the volume of the organic waste reduced 50-70 % of the original volume. Zheng (2004) reported that microorganisms reproduced very rapidly at early stage of composting due to the abundance of easily-degradable organic matter. For this, the rate of volume reduction is usually fast.

Acknowledgement:

The authors express their sincere thanks to Dr. Anil Pandey, course coordinator of Microbiology, Govt. Madhav Science College, Ujjain for providing laboratory facilities and constant encouragement.

References:

1. Banu, J. R., Yeom, I. T., Esakkiraj, S. , Kumar, N. and Logakanthi, S.2008“Biomangement of sago-sludge using an earthworm, *Eudrilus eugeniae*” Journal of Environmental Biology, 29 ,143-146.

2. Eklind ,Y., Beck-Friis, B., Bengtsson, S.,Ejlertsson, J., Kirchmann ,H., Mathisen, B., Nordkvist, E., Sonesson, U., Svensson, B. H. and Torstensson, L. 1997. Chemical characterization of source- separated organic household wastes. *Swedish Journal of Agricultural Researches*,27,167-178.
3. Finstein, M. S. and Morris, M. L. 1975. Microbiology of municipal of solid waste composting, *Advances in Applied Microbiology*.19,113-151.
4. Gray, K. R., Sherman , K. and Biddlestone, A. J.1971a. "A Review of Composting–Part 1. *Process Biochemistry*.6:32-36.
5. Haug, R. T. 1993. "*The Practical Handbook of Compost Engineering*. Lewis Publishers, London.
6. Mall , A.K., Dubey, A. and Prasad, S. 2005.Vermicompost: An inevitable tool of organic farming for sustainable agriculture. *Agrobios Newsletter*. 3,8:10-12.
7. Miller, F.C. 1996. Composting of Municipal Solid Waste and its Components. In: Palmisano AC, Barlaz MA, eds.Microbiology of Solid Waste. CRS Press, 115–154.
8. Moqsud,A. 2010. Composting barrel for sustainable organic waste management in Bangladesh, waste management, Er Sunil Kumar (Ed.), ISBN:
9. Narayan, R. 1993.Biodegradation of polymeric materials during composting. In: Hoitink A. J. and Keener H. M (eds.) *Science and Engineering of Composting: Design .Environmental, Microbiological and Utilization Aspects* Renaissance Publications, Worthington. 346.
10. Ndegwa, P. M. and Thompson, S. A. 2001.Integrating composting and vermicomposting in the treatment and bioconversion of biosolids. *Bioresource Technology* .76,107-112.
11. Norbu,T. 2002.Pre-treatment of municipal solid waste by windrow composting and vermicomposting. Ph.D. Dissertation, School of Environment Resource and Development, Thailand.
12. Rhyner, C. R., Schwartz, L. J., Wenger, R. B. and Kohrell, M.G.1995. *Waste Management and Resource Recovery*. CRC Lewis Publishers. Boca Raton.
13. Ryckeboer, J., Mergaert, J., Vaes, K., Klammer, S., De Clercq, D., Coosemans, J., Insam, H. and Swings, J.2003. A survey of bacteria and fungi occurring during composting and self-heating processes. *Annals of Microbiology*. 53:349-410.
14. Singh,R., Iyengar,L. and Rao, A. V. S. 1977. Effect of insecticides on fungal enzymes. *J. Gen. Appl. Microbiol*.23:119-126.
15. Singh,N.B.,Khare,A.K.,Bhargava,D. and Bhattacharya,S.2004.Effect of substrate depth on vermicomposting. *Journal-EN*. 85:16-21.
16. Suthar, S and. Singh, S.2008. Vermicomposting of domestic waste by using two epigeic earthworms (*Perionyx excavatus* and *Perionyx sansibaricus*).*Int. J. Environ. Sci. Tech*.5 ,1:99-106.
17. Tiquia, S.M. and Tam, N.F.Y. 2000a. Fate of nitrogen during composting of chicken litter. *Environ Pollut* (in press).
18. Vogt, K.A., Grier, C.C. and Vogt, D.J. 1986. Production, turnover, and nutrient dynamics of above- and belowground detritus of world forests. *Adv. Ecol. Res*. 15:303-377.
19. Zheng, G. D. and Chen, T. B. 2004. Dynamic of lead specialization in sewage sludge composting. *Journal of Water Sci. and Technol*. 50,9:75-82.

