



International Research Journal of Public Health (ISSN:2573-380X)



PREVALENCE OF GASTROINTESTINAL AND URINARY TRACT PARASITES AMONG STUDENTS OF ISAAC JASPER BORO COLLEGE OF EDUCATION SAGBAMA, BAYELSA STATE, NIGERIA

¹*LeBari Barine Gboeloh, ² Ayibadinipre Jennis Gbeghebo

¹Ignatius Ajuru University of Education, P.M.B 5047, Rumuolumeni, Port Harcourt, Nigeria.

²Isaac Jasper Boro College of Education, P.M.B 74, Sagbama, Bayelsa State.

ABSTRACT

The prevalence of gastrointestinal and urinary tract parasites among students of Isaac Jasper Boro College of Education (IJBCOE), Bayelsa State was investigated. Faecal and urine samples were collected from 273 presumably healthy students of the college. The samples were analyzed in the laboratory using sedimentation method for urine and formol-ether concentration method for the stool samples. A self-structured questionnaire was used to obtain some risk factors associated with the transmission of the parasites. Out of the 273 students investigated, 60 (22.0%), 31(11.4%) and 12(4.4%) were positive for gastrointestinal parasites, urinary tract parasites and coinfection of gastrointestinal and urinary tract parasites respectively. *Entamoeba histolytica* recorded the highest prevalence among the gastrointestinal parasites identified (62%), followed by *Entamoeba coli* (18.3%), *Giardia lamblia* (11.7%), *Blastocystis hominis* (5%), and *Enterobius vermicularis* (3.3%). Among the two urinary tract parasites identified, *Trichomonas vaginalis* was more prevalent (90.3%) while *Schistosoma haematobium* had 9.7%. More Female students (37.8%) were infected than males (25.7%) by gastrointestinal parasites. The age range prevalence for gastrointestinal parasites recorded 45%, 37.3%, 28.3% and 25% for 26 - 30 years, 16 – 20 years, 21 – 25 years, and 31 – 35 years respectively. Lack of awareness, consumption of faecal matter-contaminated food and water, poor sanitation, defaecation into drinking water sources were identified as risk factors for gastrointestinal parasites while engagement in unprotected sex and multiple sexual partners were recorded as risk factors for the transmission of *Trichomonas vaginalis*. Increased personal hygiene and sanitation, provision of toilet facilities and health education will mitigate the prevalence of the parasites in the study area.

Keywords: Prevalence, Gastrointestinal Parasites, Urinary tract parasites, IJBCOE, Bayelsa State.

*Correspondence to Author:

LeBari Barine Gboeloh
Ignatius Ajuru University of Education, P.M.B 5047, Rumuolumeni, Port Harcourt, Nigeria.

How to cite this article:

LeBari Barine Gboeloh, Jennis Ayibadinipre Gbeghebo. PREVALENCE OF GASTROINTESTINAL AND URINARY TRACT PARASITES AMONG STUDENTS OF ISAAC JASPER BORO COLLEGE OF EDUCATION SAGBAMA, BAYELSA STATE, NIGERIA. International Research Journal of Public Health, 2021; 5:51.

 eSciPub
eSciPub LLC, Houston, TX USA.
Website: <https://escipub.com/>

INTRODUCTION

Gastrointestinal parasites are mainly parasites of the gastrointestinal tract, they are typically protozoa and helminthes [1]. Records indicated that an estimated 65,000 species of protozoa occur in water and on land with over 100,000 species adopting the parasitic or symbiotic mode of interaction [2]. Gastrointestinal parasitic infections have widespread distribution and are prevalent occur in the poor regions of America, Asia and sub-Saharan Africa, where there is inadequate supply of potable water, personal hygiene and environmental cleanliness [3]. The most common gastrointestinal parasites include *Taenia* spp, *Ascaris lumbricoides*, *Strongyloides stercoralis* hookworms and *Schistosoma* spp, while protozoa include *Toxoplasma gondii*, *Giardia lamblia*, *Entamoeba histolytica* and *cryptosporidium parvum* [4]. These gastrointestinal parasites are serious public health concern and they constitute a leading cause of morbidity [5,6,7]. In 2010, about 438.9 million people were reportedly at risk of hookworm, 819.0 million and 464.6 million were also at risk of *Ascaris lumbricoides* and *Trichuris trichiura* respectively [8,9]. [3] reported that about 1.5 billion people globally are disease-ridden with soil-transmitted helminthes and that at least 218 million people need preventive treatment for schistosomiasis in 2015.

Gastrointestinal parasitic infections have been implicated in causing increased ill health among children in school and adult females during child bearing [10]. Children are commonly affected due to their susceptibility to poor nutrition. Apart from causing morbidity and mortality, gastrointestinal parasitic infections especially helminthes have been associated with a higher threat of nutritional anaemia [11], undernourishment, development deficits and poor educational achievements, change in resistance and immune impotence and poor cognitive skill [12, 13, 14].

In Nigeria, most human and animal waste are disposed into the soil and water bodies, environments that are suitable for the

development of the parasites. This influences the transmission of the infective stages of these gastrointestinal parasites [15].

The most common urinary tract parasites infecting humans include *S. haematobium*, *T. vaginalis* and *W. bancrofti* [16]. The health problems generated by their presence in human systems range from discomfort to serious and debilitating conditions [17]. Trichomoniasis, the disease caused by *T. vaginalis* is a pluralistic disease [18], mostly transmitted through unprotected sexual intercourse and it is the third most common cause of vaginal symptoms behind bacterial vaginosis and candidiasis [19]. It could also be non-sexually transmitted through contact with fomites and surfaces harbouring fluids from infected individuals including sharing of pants and other underwears with infected persons [18,20]. The disease is one of the commonest treatable sexually transmitted infections affecting public health [21], it could cause secondary diseases such as pelvic inflammatory disease in women, defective and low sperm count in men and preterm labour [22, 23, 24]. The most cases, the presentation is asymptomatic in both males and females, hence it is not usually and easily diagnosed [18]. Women are however, more vulnerable than men and an estimated 180 million females are infected globally [25].

Although numerous researches have been directed towards the determination of the prevalence of gastrointestinal and urinary tract parasites in various regions of the Nigeria [26, 27, 28,29], there is presently no published evidence on the prevalence of gastrointestinal parasites among students in Isaac Jasper Boro College of Education, Sagbama. This study therefore, will provide a published evidence to guide gastrointestinal parasites control programme officers and policy makers in the area of study.

MATERIALS AND METHODS

Study Area

This study was carried out among students of Isaac Jasper Boro College of Education, Sagbama. Sagbama lies between latitude

5.1522°N and longitude 6.1925°E. It is the headquarters of Sagbama Local Government Area of Bayelsa State with a population of about 187,146 [30] and has an area of 945km² (365sqm). The major occupation of the people includes fishing, farming with few civil servants. The college lies on and it is located in a swampy area and is prone to perennial flooding and lacks good toilet sanitary facilities, the only source of water within the school is a bore hole from which the entire population obtain water for various activities. Hence, most staff and students resort to open defecation in nearby bushes. Presently, students are living in temporary hostel accommodation.

Study Population

Data obtained from the admission office as at the time of this research indicated that the college has a population of eight hundred and sixty-five (865) students. The students are spread within six faculties.

Sample size

Determination of the size of sample was done according to the formula of [31]:

$$n = N / [1 + N(e)^2]$$

Where:

n = Size of sample

N= Studied population (865)

$$e = \frac{N}{\{1 + N(e)^2\}}$$

$$n = \frac{865}{\{1 + 865\{0.05\}^2\}}$$

$$n = \frac{865}{\{1 + 865 \times 0.0025\}}$$

$$n = \frac{865}{\{1 + 2.163\}}$$

$$n = \frac{865}{3.163}$$

$$n = 273$$

Sample collection

Well labeled sterile bottles (1 each) were distributed to presumably healthy students of the college alongside a self-structured questionnaire to obtain data on age, sex, marital status, occupation, nature of toilet system, sanitation,

personal hygiene and other risk factors and were trained collection their faeces. The retrieved samples were immediately conveyed to the General Hospital Laboratory Sagbama, for laboratory analysis.

Stool and urine analysis

The stool and urine samples were examined both macroscopically and microscopically.

Macroscopy: The stool samples were examined for colour, odour, texture, consistency (formed, soft, semi-soft and watery), presence of mucus or blood stains as well as the presence of segments or adult helminthes. While the urine samples were examined for their colour, clarity, turbidity and presence of blood cells.

Microscopic examination of stool samples: Parasitological investigation of stool samples was done using formol-ether concentration technique [32].

Microscopic examination of urine samples: Centrifugal sedimentation method by [32] was adopted to concentrate the parasites in the urine samples for microscopic investigation.

Ethical clearance

The ethical clearance for this work was approved by the Ministry of Health, Bayelsa State and the management of Isaac Jasper Boro College of Education, Sagbama, Bayelsa State. Each participant also gave a verbal/written consent.

RESULTS

Overall prevalence of gastrointestinal (GP) and urinary tract parasites (UTP), and coinfection of gastrointestinal and urinary tract parasites

The results indicated that out of the 273 students investigated for the presence of gastrointestinal and urinary parasites, 60 (22.0%), 31 (11.4%) and 12 (4.4%) were positive for gastrointestinal parasites, urinary tract parasites and coinfection of gastrointestinal and urinary tract parasites respectively. (Table 1).

Prevalence in relation to species of gastrointestinal and urinary tract parasites

The results of the examination of faecal samples revealed the presence of three genera of protozoa (*Entamoeba*, *Giardia*, *Blastocystis*) and a genus of nematode (*Enterobius*) while the urine test revealed the presence of two genera (*Schistosoma* and *Trichomonas*). The most prevalent among the gastrointestinal parasites was *Entamoeba histolytica* (62. %), followed by *Entamoeba coli* (18.3%), *Giardia lamblia* (11.7%), *Blastocystis hominis* (5%) and *Enterobius vermicularis* (3.3%). The prevalence

of *E. histolytica* was statistically significance ($p < 0.05$) compared to other parasites. The results further revealed that 31 (11.4%) students were infected with either of two species of urinary tract parasites. *Trichomonas vaginalis* recorded a statistically prevalence ($p < 0.05$) of 90.3% among the urinary tract parasites while *Schistosoma haematobium* recorded a prevalence of 9.7% among the urinary parasites identified (Table 2).

Table 1: Overall prevalence of gastrointestinal, urinary tract parasites and co-infection of gastrointestinal & urinary tract parasites (n = 273)

Nature of Infection	No. Infected (%)
Gastrointestinal Parasites	60 (22.0)
Urinary tract parasites	31 (11.4)
Co-infection of gastrointestinal & urinary tract parasites	12 (4.4)
Total	103(37.7%)

Table 2: Prevalence in relation to species of gastrointestinal and urinary tract parasites

Gastrointestinal Parasites	Prevalence (%)	Urinary Tract Parasites	Prevalence (%)
Protozoa			
<i>Entamoeba histolytica</i>	37 (62)	<i>Trichomonas vaginalis</i>	28(90.3)
<i>Entamoeba coli</i>	11(18.3)	<i>Schistosoma haematobium</i>	3(9.7)
<i>Blastocystis hominis</i>	3(5)		
<i>Giardia lamblia</i>	7(11.7)		
Nematode			
<i>Enterobius vermicularis</i>			
Total	2(3.3)		
	60()		31()

Coinfection of gastrointestinal and urinary tract parasites among students of IJBCOE

The result of the examination of faecal and urine samples further indicated that 12(4.4%) students had co-infection of gastrointestinal and urinary tract parasites. Out of the 12 co-infected

students, 5(1.8%) were co-infected with *E. histolytica* and *T. vaginalis*, 2(0.7%) were co-infected with *G. lamblia* and *T. vaginalis*, 1(0.4%), 1(0.4%), 3(1.1%) were co-infected with *E. histolytica* and *S. haematobium*, *B. hominis* and *T. vaginalis* and *E. coli* and *T. vaginalis* respectively (Table 3).

Table 3: Co-infection of gastrointestinal and urinary tract parasites among students of IJBCOE

Species of parasite	No. infected (%)
<i>E. histolytica</i> + <i>T. vaginalis</i>	5(1.8)
<i>E. histolytica</i> + <i>S. haematobium</i>	1(0.4)
<i>G. lamblia</i> + <i>T. vaginalis</i>	2(0.7)
<i>B. hominis</i> + <i>T. vaginalis</i>	1(0.4)
<i>E. coli</i> + <i>T. vaginalis</i>	3(1.1)
Total	12(4.4)

Risk factors associated with the transmission of gastrointestinal and urinary tract infections

The results of the study indicated that out of 273 students (male 101, female 172), 26 (25.7%) male and 65(37.8%) females were infected with at least one gastrointestinal or urinary tract

parasite. Students within the age range of 26 – 30years had the highest infection (45%), followed by 16-20years (37.3%), 21-25years (28.3%) and 31-35years (25%). Again, more single (unmarried students) were significantly ($p<0.05$) infected (34.1%) than married students (23.4%) (Table 4).

Table 4: Risk factors associated with the transmission of gastrointestinal and urinary tract infections

Variables	No. examined	No. infected (%)
Sex		
Male	101	26 (25.7)
Female	172	65 (37.8)
Age		
16 – 20	118	44 (37.1)
21 – 25	127	36 (28.3)
26 – 30	20	9 (45.0)
30 -35	8	2 (25.0)
Marital status		
Single	252	86 (34.1)
Married	21	5 (23.8)

DISCUSSION

Studies on the prevalence of gastrointestinal and urinary tract parasites among students of Isaac Jasper Boro College of Education

Sagbama was undertaken within March 2019 and February 2020. The study recorded an overall prevalence (22%) of gastrointestinal parasites among students in the study area. The 22% recorded in this study is lower than the

44.2% reported among abattoir workers in Port Harcourt by [28] and the 43.3% reported by [33] among students of a tertiary institution in Jos, Nigeria. It is also lower than the 63.5% and 40.0% recorded by [34] among rural and urban school children respectively in Benue State. The comparatively low prevalence reported in this study might be credited to the fact that only few of the students were exposed to risk factors such as contact with and consumption of contaminated food and water, lack of awareness, poor sanitary conditions, engagement in unprotected sex and multiple sexual relationships (for *T. vaginalis*) which could transmit these parasites.

On the other hand, urinary tract parasites recorded an overall prevalence of 11.4%. This is at variance with the 30.3% reported by [29] among students of Rivers State College of Health Science and Technology, Port Harcourt, Rivers State, Nigeria. Two species of urinary tract parasites (*T. vaginalis* and *S. haematobium*) were identified from the urine samples. *T. vaginalis* had the highest prevalence of 90.3% and was identified from female urine samples only. The 90.3% recorded in this study is at variance with the 11.3% reported by [35] and is in consonance with the 90.7% reported by [29]. While *T. vaginalis* recorded 90.3% prevalence rate, *S. haematobium* recorded a prevalence of 9.7% and was identified from male urine samples only. The 9.7% prevalence recorded in this study is greatly lower than the 44.84% reported by [36] among primary school pupils in Awgu Local Government Area of Enugu State. It is also lower than the 21.6% reported by [37] among school-aged children in some parts of Benue State, Nigeria. But it is higher than the 2.0% reported by [38]. The prevalence of *T. vaginalis* among the affected females may be attributed to poor personal hygiene, and sexual promiscuity [35]. The low prevalence of *S. haematobium* may be due to reduction in water contact activities and absence or presence of few intermediate host of the parasite [37].

The study further recorded a coinfection of 4.4%. This is slightly higher than the 3.6% coinfection recorded by [39] among school-aged children in Kurmi Local Government Area of Taraba State. Co-infection was higher between *E. histolytica* and *T. vaginalis* (1.8%) and was least between *E. histolytica* and *S. haematobium*, *B. hominis* and *T. vaginalis* which recorded a co-infection of 0.4% each. This is at variance with the report of [17] who recorded 3.3% co-infection of *E. histolytica*, *T. trichiura* and *A. lumbricoides* and 16.7% coinfection of *T. trichiura* and *A. lumbricoides* which were all gastrointestinal parasites. The co-infection observed in this study could be due to unhygienic life style, poor environmental health and other epidemiological influences that could encourage the spread of both gastrointestinal and urinary tract parasites [28].

Several gastrointestinal parasites were identified from faecal samples of the participants. *E. histolytica* (62%) was higher compared to other gastrointestinal parasites. This observation contradicts previous study by [28], in which *A. lumbricoides* showed higher prevalence of 45.6%. The result however agreed with the report of [40] in which *E. histolytica* had the highest prevalent rate of 20.6% among children below 5 years in Agasha, Benue State, Nigeria. *E. coli* recorded a prevalence of 18.3%. Its slightly high prevalence may be attributed to contact with faecal-contaminated food or drinks [28]. *G. lamblia* recorded a prevalence of 11.7%. This is higher than the 0.5% recorded by [41], and is lower than the 16% recorded by [42].

The low prevalence of *G. lamblia* recorded in this study may be owing to the fact that the prevalence of the parasite peaks at about age 11 years and there after decline in later life. Hence, few adults pass cysts of the parasite [43]. *B. hominis* and *E. vermicularis* recorded a prevalence of 5% and 3.3% respectively. *B. hominis* is a protozoan parasite which resides in the human intestine. Its infection is through contamination of faecally contaminated food or water [1].

The 3.3% prevalence of *E. vermicularis* recorded in this study is higher than the results reported by [44] who recorded 0.3% prevalence in their study. The slightly high prevalence of gastrointestinal parasites recorded in this study might be due to ignorance, poverty, consumption of contaminated food/water, poor environmental and poor hygiene, scarcity of drinkable water and unselective defaecation into sources of drinking water [34].

T. vaginalis recorded a prevalence of 90.3%. This is at variance with the 11.3% recorded by [35] among pregnant women in three Local Government Areas of Lagos State. The 90.3% prevalence of *T. vaginalis* is again higher than the report of [17] who recorded 10% prevalence among food handlers in Ilisha-Remo Ogun State, Nigeria. This high prevalence may be attributed to low personal hygiene and high frequency of sexual activity amongst the students, especially the females [35]. *S. haematobium* recorded a prevalence of 9.7% among the urinary tract parasites identified among the participants. This is greatly lower than the 21.6% recorded by [37] and higher than the 5.6% reported by [29]. The low prevalence of *S. haematobium* recorded in this study may be due to the absence the intermediate host of the parasite in the area of study.

Sex-related prevalence revealed a higher percentage in females (37.8%) than in males (25.6%). The high percentage in female is however lower than the 42.40% and 57.60% recorded by [45] in females and males respectively. The 37.8% in females recorded in this study is slightly higher than the 35.4% recorded by [44] among community school children in Khana Local Government Area of Rivers State, while the 25.6% in males recorded in this study is lower than the 34.5% recorded in their study. The slightly high prevalent rate noted in this study may be ascribed to unsanitary life style, poor socio-economic status, contact with soil, consumption of unwashed vegetables and fruits, drinking of contaminated water, contact with faeces [44]. In addition to the above, sexual activeness/multiple sexual partners could be

related to the slightly high prevalence in females as in the transmission of *T. vaginalis* [46].

Conclusion

The results of the study established a moderately high prevalence of gastrointestinal and urinary tract parasites among students of Isaac Jasper Boro College of Education, Sagbama. There need for health orientation of the students, provision of sanitary facilities and potable water in the study area.

Acknowledgement

The authors acknowledge the assistance rendered by the Dean, Students Affair and students that participated in the study.

Conflict of interest

The authors have no conflict of interest with regard to publication of this study.

REFERENCES

1. Arora, DR, Brij BA. Medical Parasitology. 3rd ed., CBS Publishers and Distributors; 2012.
2. Udensi JU, Mgbemena IC, Emek-Nwabunnia I, Ugochukwu MG, Awurum IN. (2005). Prevalence of Intestinal Parasites Among Primary School Children in three Geopolitical Zones of Imo State, Nigeria. *Sci J Publ Health*. 3(5-1):25-28.
3. World Health Organization. Geo-helminthiasis. Geneva: Aide memoire; No. 366,2017.
4. Norhayati M, Fatmah M, Yusof S, Edariah AB. Intestinal parasitic infections in man: A review. *The Med. J Malaysia*. 2003; 58: 296-305
5. Brooker S, Clements ACA, Bundy DAP. Global epidemiology, ecology and control of soil-transmitted helminth infections. *Adv parasitol*. 2006; 62:221-261.
6. Moore SR, Lima AA, Conaway MR, Schorling JB, Soares AM, Guerrant RL. Early Childhood diarrhea and helminthiasis associated with long-term linear growth faltering. *Int J Epidemiol*. 2001; 30(6): 147-64.
7. Crompton DW, Nesheim MC. Nutritional Impact of Intestinal Helminthiasis during the human Life. *Annual Rev Nutrit*. 2002; 22:35-59.
8. Nigeria Centre for Disease Control. Soil Transmitted Helminthes. 2017; 15:35-38. Available: <http://www.ncdc.gov.ng/diseases/info/s>. Retrieved June, 2020.
9. Pullan RL, Smith JL, Jasrasaria, R. Brooker, SJ. Global numbers of infection and disease

- burden of soil transmitted helminth infections in 2010. *Parasit Vect* 2014; 7, 37 (2014). <https://doi.org/10.1186/1756-3305-7-37>
10. Mbanugo JI, Abaziri OC. A Comparative Study of Intestinal Infection of Pregnant and Non-Pregnant Women in Nkpor, Anambra State, Nigeria. *Nig J Parasitol*. 2002; 23:19-26.
 11. Arene FOI. A handbook of applied parasitology. Paragraphics Printing. 1998; Pp. 30.
 12. Karz DE, Taylor DN. Parasitic Infections of the gastrointestinal tract. *Gastro intestinal clinic*. Worth Am. 2001; 30:395-815.
 13. Rodnquz-Morales AJ, Barbella RA, Case C, Arria M, Ravelo M, Perez H, Urdaneta O, Gervasio G, Rubio N, Maldonado A, Aguilera Y, Vilorio A, Blanco JJ, Colina M, Hernandez E, Araujo E, Cabaniel G, Benitez J, Rifakis P. Intestinal Parasitic Infections among Pregnant Women in Venezuela. *Infect. Dis. Obst Gynec.* 2006; 23125. doi: 10.1155/IDOG/2006/23125
 14. Udonsi JK. *Host Parasite "Arms Race" the Nigerian Theatre*. An Inaugural Lecture Delivered at University of Port Harcourt, 2002; Pp 68.
 15. Udensi JU, Opara FN. "Waste to Wealth: An Approach to Environmental Waste Management. *Inter J Env Health Hum Develop*. 2011; 12:66 -70.
 16. Kamau P, Aloo -Obudho P, Kabiru E, Ombacho K, Langat B, Mucheru O, Ileri L. Prevalence of Intestinal Parasitic Infections in Certified food-handlers working in food establishments in the city of Nairobi, Kenya. *J. Biomed. Res*. 2012; 26(2): 84-89.
 17. Akinboye DO, Abdullah AR, Awodele O, Akintunde TL, Effedua HL, Bamidele E. Prevalence of intestinal and urinary parasites among food-handlers in Ilisha-Remo, Ogun State. *Nigerian journal of parasitology*.: 2015; 36(2)113-118.
 18. Akinbo FO, Mokobia CN, Ande AB. Prevalence of trichomoniasis among pregnant women in Benin City. *Sahel Med J*. 2017; 20:67 - 71.
 19. Toth MD. Trichomonas and Infertility. *J Infect Dis*. 2013; 19:68-70.
 20. Ijeoma AO, Amara VN, Emmanuel IO. Prevalence of Trichomoniasis among Adults in Oru-East Local Government in Imo State, Nigeria. *Arch Clin Microbiol*. 2018; 9(2):79.
 21. von Glehn M, Ferreira Sa LCE, da Silva HDF, Machado ER. Prevalence of *Trichomonas vaginalis* in women of reproductive age at a family health clinic. *J Infect Dev Ctries* 2017; 1.(3): 269-276.
 22. Moodley P, Wilkinson D, Connolly C, Moodley J, Sturm AW. *Trichomonas vaginalis* is associated with pelvic inflammatory disease in women infected with human immunodeficiency virus. *Clin Infect Dis* 2002; 34:519-522.
 23. Ryu JS, Roh J, Lim YS, Seo MY, Choi Y. The secretory product of *Trichomonas vaginalis* decrease fertilizing capacity of mice sperm in vitro. *Asian J. Androl*. 2015;17:319.
 24. Mann JR, McDermott S, Gill J. Sexually transmitted infection is associated with increased risk of preterm birth in South Carolina women insured by Medicaid. *J Matern Neonatal Med*. 2010; 23:563-568.
 25. Swygard H, Sena AC, Hobbs MM, Gohen MS (2004). Trichomoniasis clinical manifestations diagnosis and management. *Sexually Transm Infec*. 2004; 80:91-95.
 26. Gboeloh LB, Ike-Ihunwo CN. Prevalence of Soil Transmitted Helminthes (STHs) Among Pupils of Community Primary Schools in Nkpor and Mgbodohia Communities in Rivers State, Nigeria. *South Asian J Parasit*. 2019; 2(2):1-10.
 27. Okike-Osisiogu FU, Nwoke BEB, Ukaga CN, Amaechi AA, Ezeigbo OR, Ike-Amadi CA. Prevalence of intestinal parasites and bacteria among school pupils in aba, Abia State. *Nigerian J Parasitol*. 2018; 39(1) 74 -77.
 28. Gboeloh LB, Elele K. Incidence of Gastrointestinal Parasites Among Workers in Major Abattoirs in Port Harcourt, Rivers State, Nigeria. *World Acad Sci Engr and Techn Inter J Med Health Sci*. 2013; 7 (11) 750-752.
 29. Gboeloh LB, Nelly OE, Ike-Ihunwo CN. Risk Factors Associated with Urinary Tract Parasitic Infections among Students of Rivers State College of Health Science and Technology, Port Harcourt, Rivers State, Nigeria. *Special J Med Res Health Sci*. 2019; 4(3) 1-9.
 30. National Population Commission. Demographic data in Nigeria. National Population Commission. 2006; Abuja.
 31. Taro Y. Statistics: An Introductory Analysis, 2nd Edition, New York: Harper and Row, 1967.
 32. Chesbrough M. (2018). District Laboratory Practice in Tropical Countries. 2nd Edition, Part1. New York: Cambridge University Press. 2018.
 33. Ejinaka OR, Obeta MU, Iwanse RI, Lote-Nwara IE, Nkop JP, Agbalaka PI, Friday PE. Prevalence of Intestinal parasites among Students of a Tertiary Institution in Jos Nigeria. *J Bacteriol Parasit*. 2019; 10(4) 1-4.
 34. Kpurkpur T, Sani A, Nongu C. (2016). Prevalence of Intestinal Parasitic Infections and their

- Association with Nutritional Status of Rural and Urban Pre-School Children in Benue State, Nigeria. *Interl J MCH and AIDS*. 2016; 5(2) 146-152
35. Alexander SI, Idowu ET, Otubanjo OA, Ajayi MB. Trichomoniasis among pregnant women in Ifako Ijaiye, Shomolu and Agege Local Government Area of Lagos State, Nigeria. *Nigerian J parasitol*. 2018; 39(1) 91-94.
36. Ngele KK, Okoye NT. Prevalence of Schistosomiasis infection among primary school pupils in Awgu primary school pupils in Awgu Local Government Area, Enugu State, Nigeria. *Nigerian J Parasitol*. 2016; 37(1). 11-15.
37. Obadiah HI, Idu ME, Omudu EA, Shenge MF, Ameh MO, Mwakyoga, A (2018). Studies on Schistosoma Haematobium infection in School-ages children in some parts of Benue State, Nigeria. *Nigerian J Parasitol*. 2018; 39(1).48 -51.
38. Ugochukwu O, Onwulin C, Osuala F, Dozie I, Opara F, Nwengi U. Endemicity of *Schistosomiasis* in some parts of Anambra State, Nigeria *J Med Lab and Diag*. 2013; 4:54-61.
39. Akwa YV, Jafaaru A, Elkanah OS, Chintem DGW, Njilmah JA, Thomas D, Tese T. (2018). Prevalence of gastrointestinal helminthes infections among school-aged children in Kurmi Local Government Area, Taraba State, Nigeria. *Nigerian J parasitol*. 2018; 39(2) 167 -171.
40. Utume LN, Umeh EU, Onekututu A, Omudu EA. Intestinal protozoan and helminthic diarrheal infections in children under five years old in Agasha, Benue State, north-central Nigeria. *Nigerian J Parasitol*. 2015; 36(2) 119-124.
41. Okike-Osisiogu FU, Nwoke BEB, Ukaga CN, Amaechi AA, Ezeigbo OR, Ike-Amadi CA. Prevalence of Intestinal parasites and bacteria among school pupils in Aba, Abia State. *Nigerian J Parasitol*. 2018; 39(1) 74-
42. Orji NM, Yufanyi EAC, Anene CC. Comparative Study of variability of giardiasis prevalence caused by anthropogenic activities in Fundong Health District, north-west region of Cameroon. *Nigerian Journal of Parasitology*. 2018; 39(2) 161-165.
43. Nwoke BEB. *Public Health Parasitology*. Owerri: Milestone Publishers Limited, 2018.
44. Eze CN, Owhoeli O, Ganale SS (2016). Assessment of intestinal helminthes in community school children of Khana Local Government Area, Rivers State, *Nigerian J Parasitol*. 37(1) 117-119.
45. Abah AE, Arene FOI. (2016). Intestinal parasitic infections in three geographical zones of Rivers State, Nigeria. *Nigerian J Parasitol*. 2016; 37(1) 83-85.
46. Kashibu E, Victor O, Ojumah I, Akafa R, Rikwentshe E. *Trichomonan vaginalis* infection: Prevalence and risk factors among antenatal attendees in a tertiary facility in Taraba State, Nigeria. *Nigerian J Parasitol*. 2018; 39(2) 199 -202



-
- Title: International Research Journal of Public Health
 - ISSN: 2573-380X
 - DOI: 10.28933/IRJPH
 - IF: 1.36 (citefactor)
 - Email: IRJPH@escipub.com
 - TEL: +1-281-656-1158



About the journal

The journal is hosted by eSciPub LLC. Our aim is to provide a platform that encourages publication of the most recent research and reviews for authors from all countries.

About the publisher

eSciPub LLC is a publisher to support Open Access initiative located in Houston, Texas, USA. It is a member of the largest community of professional publishers in the United States: the Independent Book Publishers Association. It hosts more than 100 Open Access journals in Medicine, Business & Economics, Agriculture, Biological Sciences, Chemistry, Education, Physical Sciences, Sociology, and Engineering and Technology.

Rapid Response Team

Please feel free to contact our rapid response team if you have any questions. Our customer representative will answer your questions shortly.



This work and its PDF file(s) are licensed under a Creative Commons Attribution-NonCommercial 4.0 International License.

Terms of Use/Privacy Policy/ Disclaimer/ Other Policies:

You agree that by using our site, you have read, understood, and agreed to be bound by all of our terms of use/privacy policy/ disclaimer/ other policies (click here for details). This site cannot and does not contain professional advice. The information on this site is provided for general informational and educational purposes only and is not a substitute for professional advice. Accordingly, before taking any actions based upon such information, we encourage you to consult with the appropriate professionals. We do not provide any kind of professional advice. The use or reliance of any information contained on this site or our mobile application is solely at your own risk. Under no circumstance shall we have any liability to you for any loss or damage of any kind incurred as a result of the use of the site or our mobile application or reliance on any information provided on the site and our mobile application. We may publish articles without peer-review. Published articles of authors are open access. Authors hold the copyright and retain publishing rights without restrictions. Authors are solely responsible for their articles published in our journals. Publication of any information in authors' articles does not constitute an endorsement by us. We make no representation or warranty of any kind, express or implied, regarding the accuracy, adequacy, validity, reliability, availability or completeness of any information that authors provided. more.....